



United States Department of the Interior



FISH AND WILDLIFE SERVICE
Washington Fish and Wildlife Office
Eastern Washington Field Office
11103 East Montgomery Drive
Spokane, Washington 99206

In Reply Refer To:
01EWF00-2018-F-0818

Michael S. Francis
Chief, Environmental Compliance Section
Walla Walla District, Corps of Engineers
201 N. Third Avenue
Walla Walla, Washington, 99362-1876

Dear Mr. Francis:

This letter transmits the U. S. Fish and Wildlife Service's (Service) Biological Opinion on the proposed Waitsburg Levee Rehabilitation Project, located in Walla Walla County, Washington, and its effects on bull trout (*Salvelinus confluentus*). Formal consultation on the proposed action was conducted in accordance with section 7 of the Endangered Species Act (Act) of 1973, as amended (16 U.S.C. 1531 *et seq.*). Your March 12, 2018, request for formal consultation was received on March 15, 2018.

The enclosed Biological Opinion is based on information provided in the March 2018, Biological Assessment, telephone conversations, emails, and other sources of information cited in the Biological Opinion. A complete record of this consultation is on file at the Eastern Washington Field Office in Spokane, Washington.


The Biological Assessment also included a request for Service concurrence with a “not likely to adversely affect” determination for bull trout critical habitat. The enclosed document includes a section separate from the Biological Opinion that addresses your concurrence request. The rationale for these concurrences is included in the concurrence section.

Michael S. Francis

2

If you have any questions regarding the enclosed Biological Opinion, our response to your concurrence request, or our shared responsibilities under the Act, please contact Michelle Eames at (509) 893-8010 or michelle_eames@fws.gov.

Sincerely,



^{for} Eric V. Rickerson, State Supervisor
Washington Fish and Wildlife Office

Enclosure

COE, Walla Walla, WA (J.Hook)

Endangered Species Act - Section 7 Consultation

BIOLOGICAL OPINION

U.S. Fish and Wildlife Service Reference:
01EWF00-2018-F-0818

Waitsburg Levee Rehabilitation Assistance Project


Walla Walla County, Washington

Federal Action Agency:

U.S. Army Corp of Engineers, Walla Walla District

Consultation Conducted By:

U.S. Fish and Wildlife Service
Washington Fish and Wildlife Office
Spokane, Washington


Eric V. Rickerson, State Supervisor
Washington Fish and Wildlife Office

5/29/18
Date

TABLE OF CONTENTS

INTRODUCTION	1
CONSULTATION HISTORY	1
CONCURRENCES.....	1
BIOLOGICAL OPINION.....	3
DESCRIPTION OF THE PROPOSED ACTION	3
ANALYTICAL FRAMEWORK FOR THE JEOPARDY AND ADVERSE MODIFICATION DETERMINATIONS	5
STATUS OF THE SPECIES: Bull Trout.....	6
ENVIRONMENTAL BASELINE: Bull Trout.....	6
EFFECTS OF THE ACTION: Bull Trout.....	11
CUMULATIVE EFFECTS: Bull Trout.....	13
INTEGRATION AND SYNTHESIS OF EFFECTS: Bull Trout.....	14
CONCLUSION: Bull Trout.....	15
INCIDENTAL TAKE STATEMENT.....	15
AMOUNT OR EXTENT OF TAKE	16
EFFECT OF THE TAKE.....	16
REASONABLE AND PRUDENT MEASURES.....	16
TERMS AND CONDITIONS	16
CONSERVATION RECOMMENDATIONS.....	17
REINITIATION NOTICE	17
LITERATURE CITED	19

APPENDICES

APPENDIX A STATUS OF THE SPECIES: BULL TROUT

FIGURES

Figure 1. Silt bluff and action area in Waitsburg, Washington. The yellow line represents a potential installation of silt fencing (Hook, J. in litt. 2018a).	3
---	---

TABLES

Table 1. Baseline aquatic habitat conditions and function.	7
---	---

ACRONYMS AND ABBREVIATIONS

BA	Biological Assessment
Service	U.S. Fish and Wildlife Service
DPS	Distinct Population Segment
CHRU	Columbia Headwaters Recovery Unit
ESA	Endangered Species Act
FMO	Foraging, Migration and Overwintering
IGDO	Inter-Gravel Dissolved Oxygen
IPCC	Intergovernmental Panel on Climate Change
ISAB	Independent Scientific Advisory Board
LPO	Lake Pend Oreille
OHWM	Ordinary High Water Mark
Opinion	Biological Opinion
PBF	Physical or Biological Features
PCE	Primary Constituent Element
PCP	Pollution Control Plan
PIT	Passive Integrated Transponder
RUIP	Recovery Unit Implementation Plan
RU	Mid-Columbia Recovery Unit
Service	U.S. Fish and Wildlife Service
SR	Spawning and Rearing
USFWS	U.S. Fish and Wildlife Service
WDFW	Washington Department of Fish and Wildlife
WWF	World Wildlife Fund

INTRODUCTION

This document represents the U. S. Fish and Wildlife Service's (Service) Biological Opinion (Opinion) based on our review of the proposed Waitsburg Levee Rehabilitation Assistance Project located in Walla Walla County, Washington, and its effects on bull trout (*Salvelinus confluentus*), in accordance with section 7 of the Endangered Species Act of 1973, as amended (16 U.S.C. 1531 *et seq.*) (Act). Your March 12, 2018, request for formal consultation was received on March 15, 2018.

This Opinion is based on information provided in the March 2018 Biological Assessment (BA), telephone conversations, emails, and other sources of information as detailed below. A complete record of this consultation is on file at the Eastern Washington Field Office in Spokane, Washington.

Your BA includes "not likely to adversely affect" determination for designated critical habitat for the bull trout, this is addressed in a concurrence section below. Your BA also included a "no effect" determination for yellow-billed cuckoo (*Coccyzus americanus*). There is no requirement for Service concurrence on "no effect" determinations. Therefore, your determinations rest with the action agency.

CONSULTATION HISTORY

The following is a summary of important events associated with this consultation:

- The Biological Assessment was received on March 12, 2018.
- Michelle Eames discussed the action with John Hook by telephone on April 24, 2018, and provided an email with questions and clarifications on April 26, 2018. We received answers to the email on the same day.
- Formal consultation was initiated on April 26, 2018.

CONCURRENCES

The Corps concluded that there would be "no effect" to yellow-billed cuckoo (*Coccyzus americanus*). There is no requirement for concurrence by the Service on "no effect" determinations. Therefore, the determination rests with the action agency.

The Proposed Action is described in detail below. It involves repair of a short portion of the levee along the Touchet River in Waitsburg Washington, through installation of a silt fence, pulling back a portion of the levee, adding rip rap, and replacing the levee to original contours. The Corps determined that the Proposed Action is not likely to adversely affect designated critical habitat for the bull trout.

Past designations of critical habitat have used the terms "primary constituent elements" (PCEs), "physical or biological features" (PBFs) or "essential features" to characterize the key

components of critical habitat that provide for the conservation of the listed species. The new critical habitat regulations (79 FR 27066) discontinue use of the terms “PCEs” or “essential features,” and rely exclusively on use of the term “PBFs” for that purpose because that term is contained in the statute. However, the shift in terminology does not change the approach used in conducting a “destruction or adverse modification” analysis, which is the same regardless of whether the original designation identified PCEs, PBFs or essential features. For those reasons, in this biological opinion, references to PCEs or essential features should be viewed as synonymous with PBFs. All of these terms characterize the key components of critical habitat that provide for the conservation of the listed species.

Several PBFs are not functional or present within the action area, and/or would not be affected by the proposed action. Effects to migration corridors (PBF #2), abundant food base (PBF #3), and water quality and quantity (PBF#8) are discussed below.

The proposed action area does not provide migration habitat during the work window in a typical water year due to low flows and high temperatures (Hook, J. *in litt.* 2018a). Mean daily stream temperature at the project site is above 20°C during the proposed work period. Additionally, mean daily discharge for the most recent available 10 year period (1980-1989) at the project site drops to 52 cfs by July 15. In a typical year, adult bull trout would have completed their migration upstream by the start of the work window.

An Environmental Protection Agency assessment (Wiseman et al 2010; Hook, J. *in litt.* 2018a) indicates that the Touchet River at the action area is characterized by an abundance of fine sediments and contains only a low abundance of sediment tolerant macroinvertebrates. The poor prey base, high stream temperatures, and poor channel complexity in the leveed portion of the river create poor conditions for forage fish during the proposed work window. As such, PBF 3 is expected to be highly impaired, and even with the proposed action would maintain the baseline condition.

The proposed project would result in short-term, localized increases in background turbidity as a result of excavation and construction of the toe wall (BA p.32). Through the use of a sediment containment barrier, sediments would be contained and allowed to settle within the action area. It is anticipated that any project related increases in background turbidity will be very limited and highly localized, and would result in insignificant effects to PBF #8.

In summary, effects from construction activities will not permanently modify existing functional PBFs associated with food base (PBF #3), migration corridors (PBF #2), or water quality and quantity (PBF #8). Therefore, we believe the Proposed Action will have no measureable effect on designated critical habitat and will be insignificant. The Service concurs that the Proposed Action is not likely to adversely affect designated critical habitat for the bull trout.

BIOLOGICAL OPINION

DESCRIPTION OF THE PROPOSED ACTION

A federal action means all activities or programs of any kind authorized, funded, or carried out, in whole or in part, by federal agencies in the United States or upon the high seas (50 CFR 402.02). The Proposed Action is within the mainstem Touchet River, a major tributary to the Walla Walla River (Fig.1).




Figure 1. Silt bluff and action area in Waitsburg, Washington. The yellow line represents a potential installation of silt fencing (Hook, J. in litt. 2018a).

A silt bluff is eroding on private land upstream of the levee. The Corps proposes to construct a riprap buttress at the upstream end of the levee to armor and protect the levee tie-in point so that high water does not go around the levee. Equipment to be used could include an excavator, dump trucks, cranes, or similar machinery. To construct the buttress the levee would be excavated and the bench surface on top of the levee would be extended upstream to provide a stable surface to stage excavation machinery. Once a work platform was stabilized, the levee would be excavated to approximately 20 feet back from its current slope. A toe wall would be constructed at the base of the levee and large riprap would be placed to construct a robust revetment at the tie in point of the levee. The riprap would then be filled with smaller rock and sand, and the original contours

of the levee restored. The levee surface would then be reseeded (BA p.9). Excavation would take place along less than 55 linear feet of shoreline (Hook, J. *in litt.* 2018c)

To minimize any further impingement of the river at high flows, the riprap buttress would not extend into the river. Riprap would be scavenged if available, but it is expected that riprap will need to be brought in from off site. Imported riprap would be clean quarry rock up to 4 foot in width. Construction work would be performed from the top of the levee or from the yard located behind the levee. Construction equipment would be staged at a secured area in Waitsburg, not along the Touchet River. Material excavated from the levee would be trucked to an offsite disposal location.

All work would be conducted during the normal in-water work window of July 15 to August 15, 2018. The work window is 4 weeks, but within that period the entirety of the work is expected to take 4 to 5 days at most (Hook, J. *in litt.* 2018a). Excavation could occur on any of those days.

Conservation Measures

The Corps proposes the following conservation measures as part of the proposed action.

1. All heavy equipment (i.e., crane and excavator, dump trucks) will access the project site via existing roadways, parking areas, the top of the levee, and disturbed upland areas.
2. A Pollution Control Plan (PCP) will be prepared by the Contractor and carried out commensurate with the scope of the project that includes the following:
 - BMPs to confine, remove, and dispose of construction waste.
 - Procedures to contain and control a spill of any hazardous material.
 - Steps to cease work under high flow conditions.
3. Only enough supplies and equipment to complete the project will be stored at the staging area and only the material and equipment required for each day's work will be on the actual work site (clarified in Hook, J. *in litt.* 2018a).
4. All equipment will be inspected daily for fluid leaks, any leaks detected will be repaired before operation is resumed.
5. Before operations begin, and as often as necessary during operation, all equipment that will be used below the OHWM will be steam cleaned until all visible oil, grease, mud, and other visible contaminants are removed.
6. Stationary power equipment operated within 150 feet of the Touchet River will be diapered to prevent leaks.
7. A sediment containment barrier or silt fence would be used to contain sediment within the proposed action area (BA p.10). To construct the buttress the Corp will first use seines or block nets to "push" any fish near the left bank waterward into deeper flow of the river and isolate an area along the left bank approximately 10-feet into the channel from the toe of the previous/existing levee (Driscoll, D. *in litt.* 2018). Workers will create a temporary sediment barrier within the isolated area using sandbags, straw bales and plastic. Using only dipnets, no electrofishing, fish will be salvaged from the isolated area and it will be allowed to dewater naturally or water will be pumped to an upslope location to infiltrate back into the river. The fencing would be left in place after completion of the work until sediments had settled behind it. Turbidity releases would then be limited to sediment disturbance from installation and removal of the curtain (Hook, J. *in litt.* 2018a).

Action Area

The action area is defined as all areas to be affected directly or indirectly by the federal action and not merely the immediate area involved in the action (50 CFR 402.02). In delineating the action area, we evaluated the farthest reaching physical, chemical, and biotic effects of the action on the environment.

The action area for this proposed Federal action is based on the geographic extent of turbidity, and the extent of the silt fence and the excavation area. Therefore, the action area includes the Touchet River from the upstream edge of the action, to approximately 200 feet downstream of the end of the silt fence (a rough estimate of the extent of the turbidity plume). The action area includes the levee excavation area, and extends landward about 20 feet from the toe of the levee, and water-ward to the extent of the silt fence and excavation, about 10 feet in width (Hook, J. *in litt.* 2018b).

ANALYTICAL FRAMEWORK FOR THE JEOPARDY AND ADVERSE MODIFICATION DETERMINATIONS

Jeopardy Determination

The following analysis relies on the following four components: (1) the *Status of the Species*, which evaluates the rangewide condition of the listed species addressed, the factors responsible for that condition, and the species' survival and recovery needs; (2) the *Environmental Baseline*, which evaluates the condition of the species in the action area, the factors responsible for that condition, and the relationship of the action area to the survival and recovery of the species; (3) the *Effects of the Action*, which determines the direct and indirect impacts of the proposed Federal action and the effects of any interrelated or interdependent activities on the species; and (4) *Cumulative Effects*, which evaluates the effects of future, non-federal activities in the action area on the species.

In accordance with policy and regulation, the jeopardy determination is made by evaluating the effects of the proposed federal action in the context of the species' current status, taking into account any cumulative effects, to determine if implementation of the proposed action is likely to cause an appreciable reduction in the likelihood of both the survival and recovery of listed species in the wild.

The jeopardy analysis in this Opinion emphasizes the rangewide survival and recovery needs of the listed species and the role of the action area in providing for those needs. It is within this context that we evaluate the significance of the proposed Federal action, taken together with cumulative effects, for purposes of making the jeopardy determination.

STATUS OF THE SPECIES: Bull Trout

The bull trout was listed as a threatened species in the coterminous United States in 1999. Throughout its range, bull trout are threatened by the combined effects of habitat degradation, fragmentation, and alterations associated with dewatering, road construction and maintenance, mining, grazing, the blockage of migratory corridors by dams or other diversion structures, poor water quality, incidental angler harvest, entrainment, and introduced non-native species (64 FR 58910 [Nov. 1, 1999]). Since the listing of bull trout, there has been very little change in the general distribution of bull trout in the coterminous United States, and we are not aware that any known, occupied bull trout core areas have been extirpated (USFWS 2015a, p. iii).

The 2015 recovery plan for bull trout identifies six proposed recovery units within the listed range of the species (USFWS 2015a, p. 33). Each of the recovery units are further organized into multiple bull trout core areas, which are mapped as non-overlapping watershed-based polygons, and each core area includes one or more local populations. Within the coterminous United States we currently recognize 109 occupied core areas, which comprise 600 or more local populations of bull trout (USFWS 2015a, p. 34). Core areas are functionally similar to bull trout metapopulations, in that bull trout within a core area are much more likely to interact, both spatially and temporally, than are bull trout from separate core areas.

The Service has also identified a number of marine or mainstem riverine habitat areas outside of bull trout core areas that provide foraging, migration, and overwintering (FMO) habitat that may be shared by bull trout originating from multiple core areas. These shared FMO areas support the viability of bull trout populations by contributing to successful overwintering survival and dispersal among core areas (USFWS 2015a, p. 35).

For a detailed account of bull trout biology, life history, threats, demography, and conservation needs, refer to Appendix A: Status of the Species - Bull Trout.

ENVIRONMENTAL BASELINE: Bull Trout

Regulations implementing the Act (50 CFR 402.02) define the environmental baseline as the past and present impacts of all Federal, State, or private actions and other human activities in the action area. Also included in the environmental baseline are the anticipated impacts of all proposed federal projects in the action area that have undergone section 7 consultation, and the impacts of state and private actions which are contemporaneous with the consultation in progress.

Current Condition of the Bull Trout in the Action Area

This section analyzes the current condition of the bull trout in the action area, the factors responsible for that condition, and the intended role of the action area in the conservation of the Mid-Columbia recovery unit for the bull trout. The action area lies within the Columbia River Distinct Population Segment which is identified in the bull trout listing rule for the coterminous United States (64 FR 28910). The condition of the species and critical habitat in the action area

described below uses data compiled and summarized during the comprehensive Bull Trout 5-Year Review process (USFWS 2008, USFWS 2015b), the draft bull trout Recovery Plan chapters (USFWS 2002, USFWS 2004), the final Recovery Unit Implementation Plan for the Mid-Columbia River (USFWS 2015c), the Final Rule designating revised bull trout critical habitat (75 FR 63898), and other available local information.

The Touchet River, a tributary to the Walla Walla River, is located within the Touchet River Core Area, which is located in the Mid-Columbia Recovery Unit. The Touchet River drains the northern and northwestern portions of the Walla Walla Basin before entering the lower mainstem Walla Walla River about 21.6 miles upstream of the Columbia River near the community of Touchet, Washington. The North Fork, South Fork, and Wolf Fork feed into the Touchet River at the base of the Blue Mountains near the City of Dayton. Lewis Creek and Spangler Creek are main tributaries to the North Fork Touchet River, while the Burnt Fork is the main tributary to the South Fork Touchet River.

Factors Responsible for the Condition of the Species

Table 1 summarizes our assessment of aquatic habitat function using the Matrix of Diagnostics/Pathways and Indicators (USFWS 1998). The matrix is a tool for describing whether habitat is “properly functioning,” “functioning at risk,” or “not properly functioning” at the scales of the action area and watershed.

Table 1. Baseline aquatic habitat conditions and function.

	Action Area	Watershed
Water Quality		
Temperature	Functioning At Risk	Functioning At Risk
Sediment	Functioning At Risk	Functioning At Risk
Chemical Contamination & Nutrients	Functioning Appropriately	Functioning Appropriately
Habitat Access		
Physical Barriers	Functioning at Unacceptable Risk	Functioning at Unacceptable Risk
Habitat Elements		
Substrate	Functioning At Risk	Functioning At Risk
Pool Frequency	Functioning At Risk	Functioning At Risk
Large Woody Debris	Functioning At Risk	Functioning At Risk
Riparian Habitat	Functioning At Risk	Functioning At Risk
Channel Conditions & Dynamics		
Streambank Condition	Functioning At Risk	Functioning At Risk
Floodplain Connectivity	Functioning at Unacceptable Risk	Functioning At Risk
Flow Hydrology		
Change in Peak/Base Flows	Functioning At Risk	Functioning At Risk
Drainage Network Increase	Functioning At Risk	Functioning At Risk

Current conditions within the middle section of the Touchet River where the action area is located are severely degraded relative to historic conditions: a reflection of a multitude of actions whose effects frame the environmental baseline in the action area. Current conditions in the action area are degraded relative to historic conditions primarily because of the almost basin-wide confinement of the river within levee structures created for road, agriculture and rural development. Vegetation along the river in the action area is limited in extent, with some patches of shrubs, and open areas, including yards. In most of the action area development (rural, residential and agriculture) extends to the river banks. The largest irrigation diversions in the river are located downstream of the action area and they are a primary contributor to low water or dewatered conditions in the lower river from early summer to late fall.

The Touchet River in the action area is located within the City of Waitsburg, Washington. The left (south) bank of the river upstream of the action area is composed of flood risk reduction levees constructed primarily of riprap with isolated pockets of shrubby vegetation that is frequently removed. Immediately behind the levee in the action area are business and residential structures. The Highway 12 bridge is just upstream of the action area.

Elevated water temperatures from factors such as damaged riparian vegetation, increased sedimentation, and decreased water flows have reduced habitat quality for bull trout in the Touchet drainage (Mendel et al. 2007). Introduced brown trout and rainbow trout likely compete with native bull trout for food and habitat, while introduced non-native walleye and small mouth bass in the lower reaches of the Touchet and mainstem Walla Walla River pose a predatory risk to juveniles and sub-adults in the basin. There are a few partial or seasonal barriers to movement in the core area that limit connectivity between local populations. Flood control levees have confined the river and reduced channel complexity and wood recruitment.

Water temperatures in summer are high. The nearest monitoring, at Bolles bridge approximately 4.2 river miles downstream, indicates that mean daily temperatures in the main stem Touchet River are consistently above 20 °C during the summer work window (Figure 10 in BA p.20). While these temperatures are not immediately lethal for bull trout, they are well outside the preferred temperature range. Additionally, migratory patterns would suggest that few bull trout would be present in the lower Touchet River during the proposed work window.

Overall, rural development, agricultural practices, channelization, and road building have degraded the environmental baseline of the action area. Thus, many aquatic habitat indicators are not properly functioning in the action area.

Conservation Role of the Action Area

Historically, bull trout were thought to be widely distributed in the Touchet River watershed (Mendel et al. 2003). Currently, local populations in the Touchet River core area occur in the North Fork, Wolf Fork, and in the Burnt Fork of the South Fork Touchet River (Kassler and Mendel 2007). Both fluvial migratory and resident forms are present throughout. However, recent telemetry and PIT tag data indicate migratory bull trout in the Touchet River core remain within the overall Walla Walla basin, foraging and overwintering in the lower Touchet drainage or mainstem Walla Walla River, and do not migrate further downstream into the Columbia River

(Schaller et al. 2014). Kassler and Mendel (2007) determined that more than 50 percent of migratory bull trout in the Touchet River core area originate from the Wolf Fork population. Spawning also occurs in Spangler and Lewis Creeks; however, genetics from individuals from each tributary were not distinguishable from either North Fork or Wolf Fork individuals (Kassler and Mendel 2007). Redd counts in the North Fork and Wolf Fork between 1999 and 2013 suggest that these two local populations are stable (Mendel et al. 2014). However, redd count data for the Burnt Fork of the South Fork Touchet is more limited. Bull trout redds were first observed in 2000, but not detected in 2003 and 2004 (Mendel et al. 2004 and Mendel et al. 2009). Since 2005, access to complete surveys in the Burnt Fork has been restricted across private property (Mendel et al. 2014).

Currently, the lower Touchet River in the action area provides FMO for bull trout. The FMO habitat is important to bull trout of the Mid-Columbia Recovery Unit for maintaining diversity of life history forms and for providing access to productive foraging areas. Many bull trout of the Touchet River core area are fluvial or resident, and therefore rely on middle portions of the basin for migrating, overwintering, extended rearing and growth to maturity (USFWS 2015c). However, the action area does not provide suitable bull trout spawning habitat and provides little or no functional rearing habitat. The primary threats identified in the 2015 Recovery Plan include Upland/Riparian Land Management, Instream Impacts, Water Quality, Connectivity Impairments, and Nonnative Fishes (USFWS 2015c, p. C-12). Most identified threats are generally specific to the core area as a whole; however some do occur in the Touchet River action area.

Climate Change

Consistent with Service policy, our analyses under the ESA include consideration of ongoing and projected changes in climate. The term “climate” refers to the mean and variability of different types of weather conditions over time, with 30 years being a typical period for such measurements, although shorter or longer periods also may be used (IPCC 2014a, pp. 119-120). The term “climate change” thus refers to a change in the mean or variability of one or more measures of climate (e.g., temperature or precipitation) that persists for an extended period, typically decades or longer, whether the change is due to natural variability, human activity, or both (IPCC 2014a, p. 119). Various types of changes in climate can have direct or indirect effects on species and critical habitats. These effects may be positive, neutral, or negative, and they may change over time. The nature of the effect depends on the species’ life history, the magnitude and speed of climate change, and other relevant considerations, such as the effects of interactions of climate with other variables (e.g., habitat fragmentation) (IPCC 2014b, pp. 64, 67-69, 94, 299). In our analyses, we use our expert judgment to weigh relevant information, including uncertainty, in our consideration of various aspects of climate change and its effects on species and their critical habitats. We focus in particular on how climate change affects the capability of species to successfully complete their life cycles, and the capability of critical habitats to support that outcome. Our analyses under the Act include consideration of ongoing and projected changes in climate. The terms “climate” and “climate change” are defined by the Intergovernmental Panel on Climate Change (IPCC). The term “climate” refers to the mean and variability of different types of weather conditions over time, with 30 years being a typical period for such measurements, although shorter or longer periods also may be used (IPCC 2007a,

p. 78). The term “climate change” thus refers to a change in the mean or variability of one or more measures of climate (e.g., temperature or precipitation) that persists for an extended period, typically decades or longer, whether the change is due to natural variability, human activity, or both (IPCC 2007a, p. 78).

In the Pacific Northwest, most models project warmer air temperatures, increases in winter precipitation, and decreases in summer precipitation. Warmer temperatures will lead to more precipitation falling as rain rather than snow. As the seasonal amount of snow pack diminishes, the timing and volume of stream flow are likely to change and peak river flows are likely to increase in affected areas. Higher air temperatures are also likely to increase water temperatures (Independent Scientific Advisory Board (ISAB) 2007).

All life stages of the bull trout rely on cold water. Increasing air temperatures are likely to impact the availability of suitable cold-water habitat. For example, ground water temperature is generally correlated with mean annual air temperature, and has been shown to strongly influence the distribution of many char species. Ground water temperature is linked to bull trout selection of spawning sites, and has been shown to influence the survival of embryos and early juvenile rearing of bull trout (Rieman et al. 2007). Increases in air temperature are likely to be reflected in increases in both surface and groundwater temperatures.

Migratory bull trout can be found in lakes, large rivers, and marine waters. Effects of climate change on lakes are likely to impact migratory adfluvial bull trout that seasonally rely upon lakes for their greater availability of prey and access to tributaries. Climate-warming impacts to lakes will likely lead to longer periods of thermal stratification, and cold-water fish may be restricted to the middle and bottom portions of the water column for longer periods of time in the summer. Deeper thermoclines resulting from climate change may further reduce the area of suitable temperatures and intensify competition for food (WWF 2003).

Bull trout require very cold water for spawning and incubation. Suitable spawning habitat is often found in accessible higher elevation tributaries and headwaters of rivers. However, impacts on hydrology associated with climate change are related to shifts in timing, magnitude and distribution of peak flows that are also likely to be most pronounced in these high elevation stream basins (Battin et al. 2007). The increased magnitude of winter peak flows in high elevation areas is likely to impact the location, timing, and success of spawning and incubation for the bull trout and Pacific salmon species. Although lower elevation river reaches are not expected to experience as severe an impact from alterations in stream hydrology, they are unlikely to provide suitably cold temperatures for bull trout spawning, incubation, and juvenile rearing. As climate change progresses and stream temperatures warm, thermal refugia will be critical to the persistence of many bull trout populations.

There is still a great deal of uncertainty associated with predictions relative to the timing, location, and magnitude of future climate change. It is also likely that the intensity of effects will vary by region (ISAB 2007) although the scale of that variation may exceed that of States. For example, several studies indicate that climate change has the potential to impact ecosystems in nearly all streams throughout the State of Washington (ISAB 2007, Battin et al. 2007, Rieman et al. 2007). In streams and rivers with temperatures approaching or at the upper limit of allowable

water temperatures, there is little if any likelihood that bull trout will be able to adapt to or avoid the effects of climate change/warming. There is little doubt that climate change is and will be an important factor affecting bull trout distribution. As its distribution contracts, patch size decreases and connectivity is truncated, bull trout populations that may be currently connected may face increasing isolation, which could accelerate the rate of local extinction beyond that resulting from changes in stream temperature alone (Rieman et al. 2007). Due to variations in landform and geographic location across the range of the bull trout, it appears that some populations face higher risks than others. Bull trout in areas with currently degraded water temperatures and/or at the southern edge of its range may already be at risk of adverse impacts from current as well as future climate change.

Recent climate change modeling indicates that the Touchet drainage is at high risk for reduced instream flows, elevated water temperatures, and reduced habitat suitability into the future and existing habitat threats will likely be exacerbated (Schaller et al. 2014).

EFFECTS OF THE ACTION: Bull Trout

The effects of the action refers to the direct and indirect effects of an action on the species or critical habitat, together with the effects of other activities that are interrelated or interdependent with that action, that will be added to the environmental baseline (50 CFR 402.02). Indirect effects are those that are caused by the proposed action and are later in time, but still are reasonably certain to occur.

General effects that will occur because of the proposed actions are: (1) turbidity caused by in-channel work including placement and removal of the silt fencing, (2) temporary loss of small amount of forage area and alteration of substrates during the project, and (3) injury or death from entrainment during excavation. Indirect effects include behavioral changes resulting from a short period of elevated turbidity levels during channel re-watering following construction. Effects to bull trout will be minimized through a limited summer work window of July 15 through August 15. Because of the low number of bull trout present in the Touchet River near the action area during the work window the risk of harming a listed species during construction is low, though not discountable (BA p.30).

Water quality and Turbidity

It is difficult to give an estimate of the length and duration of a turbidity pulse from installation and removal of the silt fencing (Hook, J. *in litt.* 2018a), however for similar projects the Service has estimated 200 feet. While flow in the Touchet River will be quite low during the work period, sediments in the action area are relatively fine and prone to suspension. Armoring and restoring the levee to its original contours would reduce sediment load and turbidity during annual high spring flows in the long term.

Short-term, localized project-related increases in background turbidity levels will likely occur as a result of proposed excavation activities and during the establishment of the toe wall (BA p.31). In the short term, increases in turbidity can reduce forage quantity for salmonids, and disrupt behavioral patterns such as feeding and sheltering. Exposure duration is a critical determinant of physical or behavioral turbidity effects. Salmonids have evolved in systems that periodically

experience short-term pulses (days to weeks) of high suspended sediment loads, often associated with flood events, and are adapted to such seasonal high pulse exposures (NMFS 2011). The entirety of the work is expected to take four to five days at most (Hook, J. *in litt.* 2018a). Excavation could occur on any of the work days, however the silt fencing would be left in place after completion of the work until sediments had settled behind it (Hook, J. *in litt.* 2018a).

Given the existing substrate conditions (primarily sandy-silt) and highly erodible silt bluff near the action area, there is a significant risk of sedimentation and turbidity due to the proposed project activities (BA p.31). However any sediment mobilized should be contained at the action area by the silt barrier. Given that few individuals of listed species are likely to be present and that sediment would be contained by a silt barrier, it is anticipated the any project related increases in background turbidity will be very limited and highly localized. As such, short-term increases in background turbidity are not expected to result in long-term adverse effects to ESA-listed fish species, or significant net change in function of the in-stream habitat. Turbidity releases would then be limited to sediment disturbance from installation and removal of the silt fence expected to be very low (barely visible), the duration will be a matter of minutes to an hour, and downstream transport will be minimized by low flows (Newcombe and Macdonald 1991). Temporary increases in suspended sediment concentrations have highly variable effects on fish, ranging from behavioral effects including alarm reactions and avoidance responses to sub-lethal effects including reduced feeding and physiological stress (Newcombe and Jensen 1996).

Equipment operating near and over the river channel within the action area presents a potential source of chemical contamination (BA p.31). Accidental spills of construction materials or petroleum products would adversely affect water quality and potentially impact bull trout, however, development and implementation of a Pollution Control Plan (PCP) that will include containment measures and spill response for construction-related chemical hazards and implementation of other conservation measures will significantly reduce the likelihood for chemical releases within the action area, making effects discountable.

Alteration of Substrates and Effects to Foraging Habitats

The proposed project will result in the alteration of in-water substrates associated with excavation of the levee and construction of the toe wall at its base (BA p. 30-31). There will be limited excavation of the river bed, at the base of the levee, to accommodate the construction of the toe wall, estimated to be 55 feet long by 10 feet wide within the channel. Current sandy-silt substrates at the eroding levee face would be replaced with riprap at the toe wall.

In general, the environmental baseline within the project action area has been degraded by development and human activity, and provides very little habitat complexity for salmonids (BA p. 30-31). As such, given the existing baseline conditions and substrates (primarily sandy-silt), proposed timing of in-water work (outside the peak migration stages), relative size of the action area, and proposed excavation techniques, it is unlikely that the proposed alteration of existing substrates will result in long-term adverse effects to forage opportunities for bull trout. Forage quantity for juvenile fish may be temporarily reduced within the immediate in-water work area as benthic organisms become disturbed by excavation; however, recolonization of benthic

organisms will likely occur within a month following project completion (NMFS 2009 as referenced in BA), and as described in the baseline, forage quality is not high in the action area (Wiseman et al 2010; Hook, J. in litt. 2018a). However, forage species will begin to re-colonize the area after project completion via drift and migration. Given the size of the disturbed area, the amount of available local habitat, and the short-term nature of the action, the Service expects short-term (from a few days up to a couple weeks) localized reduced productivity followed by a return to pre-project conditions. In addition, bull trout densities in the area are expected to be low, and they will be able to move up or downstream to other foraging areas. The Service does not expect food availability to result in any injury or mortality or appreciably alter survival or fitness of any of those fish within the action area; therefore effects from the short term loss of food availability are insignificant.

Salvage Activities and Excavation and Entrainment

As described in the proposed action, the Corps will implement a sediment containment barrier or silt fence, as part of that they will first use seines or block nets to “push” any fish near the left bank waterward into deeper flow of the river and isolate an area along the left bank approximately 10-feet into the channel from the toe of the previous/existing levee (Driscoll, D. in litt. 2018). Using only dipnets, and no electrofishing, fish will be salvaged from the isolated area within the sediment barrier, and it will be allowed to dewater naturally or water will be pumped to an upslope location to infiltrate back into the river.

Excavation activities will directly affect instream habitat that supports bull trout and low numbers of bull trout are expected to be present at the time of construction. While work conducted below the OHWM of the Touchet River would be conducted during the approved in-water work windows, or in complete isolation from the adjacent waters, it is likely that bull trout will be present and exposed to excavation activities. Entrainment may occur if bull trout are missed during the salvage and are trapped in the bucket of the excavator during excavation of the bank at the action area and during the construction of the toe wall (BA p.30-31).

The potential for bull trout exposure during salvage or entrainment is low, given the proposed timing of in-water work (July 15 through August 15), location of proposed excavation activities (i.e., near the shoreline), use of an open bucket excavator, initial establishment of a toe wall, and relatively slow speed of excavation; however it is not discountable, especially for smaller subadult bull trout.

Based on low densities of bull trout expected in the action area during the proposed work window, the Service estimates one bull trout would be exposed to the salvage and excavation and be killed or injured. The maximum extent of the isolation area and excavation area within the river channel is 10 feet by 55 feet, for a total of 550 square feet.

CUMULATIVE EFFECTS: Bull Trout

Cumulative effects include the effects of future state, tribal, local, or private actions that are reasonably certain to occur in the action area considered in this Opinion. Future federal actions

that are unrelated to the proposed action are not considered in this section because they require separate consultation pursuant to section 7 of the Act.

The Touchet River watershed is the location of numerous habitat restoration actions taken by local agencies and watershed groups: the Confederated Tribes of the Umatilla Indian Reservation, the Snake River Salmon Recovery Board, WDFW, Service and U.S. Forest Service. We are not aware of any specific future non-federal actions that are reasonably certain to occur in the action area that are likely to contribute to cumulative effects on bull trout or designated critical habitat. Many, if not all, of the possible future actions that would or might have significance for bull trout and designated steelhead critical habitat are likely to have an independent federal nexus, and would therefore be subject to the requirements of a separate section 7 consultation process.

The Service does expect that Walla Walla County will continue to experience population growth, and that there will be additional residential, urban, agricultural and industrial development along Mill Creek. This future development may have measurable effects to floodplain and riparian functions, the hydrological regime, water quality, and instream habitat. Any potential future development would likely continue the current practices of extensive levee structures that cause measurable negative effects to floodplain and riparian functions, the hydrological regime, water quality, and instream habitat. We expect that the cumulative effects of future State, Tribal, local, and private actions are likely to maintain the current impaired conditions and contribute to continued inability of critical habitat to recover.

INTEGRATION AND SYNTHESIS OF EFFECTS: Bull Trout

The waters within the action area provide FMO habitat for bull trout. FMO habitat is important to bull trout of the Mid-Columbia Recovery Unit for maintaining diversity of life history forms and for providing access to productive foraging areas. Bull trout in the action area likely originate from either of the Touchet River local populations. Many bull trout of the Touchet River core area are fluvial or resident, and therefore rely on middle portions of the basin for migrating, overwintering, extended rearing and growth to maturity (USFWS 2015c). However, the action area does not provide suitable bull trout spawning habitat and provides little or no functional rearing habitat.

The proposed action incorporates conservation measures which will reduce effects to habitat and avoid and minimize impacts during construction. The proposed action will affect foraging and migration habitat for subadult and adult bull trout. Temporary effects will result from exposure to elevated turbidity. The proposed action will have no measurable, permanent adverse effects to water quantity or quality; water quality effects will be short in duration. With full and successful implementation of the proposed BMPs, the Service expects that only low numbers of subadult and adult bull trout will be exposed to turbidity from construction activities and suffer insignificant effects.

The herding, capture, and handling of bull trout for salvage purposes will result in direct take (kill, capture, injury). However, the direct take resulting from salvage operations will minimize

the incidental take of individual bull trout from stream diversion and dewatering activities. Direct adverse effects to bull trout are expected from excavation in the river channel at the toe of the levee. The Service expects that subadult or adult bull trout may occupy the action area at any time of year, although they are less likely during high temperature and low flow seasons of the year. Their distribution and density varies with flow. The Service expects only one bull trout will be exposed to excavation within the 550 square foot area of the silt fence.

The Service expects that the proposed action will have no measurable effect on the relative sizes of the fluvial and resident individuals contributing to the core area's local populations. The action will have no effect on distribution at the scale of the local populations or Touchet River core area. The anticipated direct and indirect effects of the action, combined with the effects of interrelated and interdependent actions, and the cumulative effects will not appreciably reduce the likelihood of survival and recovery of the species. The anticipated direct and indirect effects of the action (permanent and temporary) will not measurably reduce bull trout reproduction, numbers, or distribution at the scale of the Touchet River core area or the Mid-Columbia Recovery Unit. The anticipated direct and indirect effects of the action will not alter the status of bull trout at the scale of the coterminous range.

CONCLUSION: Bull Trout

After reviewing the current status of bull trout, the environmental baseline for the action area the effects of the Proposed Action and the cumulative effects, it is the Service's Opinion that the Action, as proposed, is not likely to jeopardize the continued existence of the bull trout.

INCIDENTAL TAKE STATEMENT

Section 9 of the Act and federal regulation pursuant to section 4(d) of the Act prohibit the take of endangered and threatened species, respectively, without special exemption. Take is defined as to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct. *Harm* is defined by the Service as an act which actually kills or injures wildlife. Such an act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavior patterns, including breeding, feeding, or sheltering (50 CFR 17.3). *Harass* is defined by the Service as an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding, or sheltering (50 CFR 17.3). Incidental take is defined as take that is incidental to, and not the purpose of, the carrying out of an otherwise lawful activity. Under the terms of section 7(b)(4) and section 7(o)(2), taking that is incidental to and not intended as part of the agency action is not considered to be prohibited taking under the Act provided that such taking is in compliance with the terms and conditions of this Incidental Take Statement.

The measures described below are non-discretionary, and must be undertaken by the Corps so that they become binding conditions of any grant or permit issued to the City, as appropriate, for

the exemption in section 7(o)(2) to apply. The Corps has a continuing duty to regulate the activity covered by this Incidental Take Statement. If the Corps 1) fails to assume and implement the terms and conditions or 2) fails to require the (applicant) to adhere to the terms and conditions of the incidental take statement through enforceable terms that are added to the permit or grant document, the protective coverage of section 7(o)(2) may lapse. In order to monitor the impact of incidental take, the Corps must report the progress of the action and its impact on the species to the Service as specified in this Incidental Take Statement [50 CFR 402.14(i)(3)].

AMOUNT OR EXTENT OF TAKE

The Service anticipates that one bull trout will be taken as a result of this proposed action. The incidental take is expected to be in the form of kill, injure, capture. The Service expects that incidental take of bull trout will be difficult to detect or quantify for the following reasons: 1) the low likelihood of finding dead or injured adults or sub-adults; 2) the ability of subadult bull trout to hide in interstitial spaces makes them difficult to find, and 3) bull trout will be unlikely to be seen in an excavator full of dirt. However, the following level of take of this species can be anticipated by loss of 550 square feet of aquatic habitat from excavation at the toe of the dike because this is where subadult bull trout could hide and be entrained during the excavation, resulting in harm.

EFFECT OF THE TAKE

In the accompanying Opinion, the Service determined that the level of anticipated take is not likely to result in jeopardy to the species or destruction or adverse modification of critical habitat.

REASONABLE AND PRUDENT MEASURES

The conservation measures negotiated in cooperation with the Corps and included as part of the proposed action constitute all of the reasonable measures necessary to minimize the impacts of incidental take. On that basis, no Reasonable and Prudent Measures except for monitoring and reporting requirements are included in this Incidental Take Statement.

- 1) Monitor and report incidental take as measured by square feet of excavation within the channel.

TERMS AND CONDITIONS

In order to be exempt from the prohibitions of section 9 of the Act, the Corps must comply with the following terms and conditions, which implement the reasonable and prudent measures, described above and outline required reporting/monitoring requirements. These terms and conditions are non-discretionary.

- 1) The Corps shall submit a monitoring report to the Eastern Washington Field Office in Spokane, Washington (Attn: Michelle Eames), by March 1 following implementation of the action. The report shall include, at a minimum, the following: (a) dates, times, and locations of excavation activities, (b) square feet of excavation within the channel, and (c) number and size of bull trout observed during the action.

The Service believes that no more than one bull trout associated with salvage and excavation will be killed or injured as a result of the Proposed Action. The reasonable and prudent measures, with their implementing terms and conditions, are designed to minimize the impact of incidental take that might otherwise result from the proposed action. If, during the course of the action, this level of incidental take is exceeded, such incidental take represents new information requiring reinitiation of consultation and review of the reasonable and prudent measures provided. The Federal agency must immediately provide an explanation of the causes of the taking and review with the Service the need for possible modification of the reasonable and prudent measures.

The Service is to be notified within three working days upon locating a dead, injured or sick endangered or threatened species specimen. Initial notification must be made to the nearest U.S. Fish and Wildlife Service Law Enforcement Office. Notification must include the date, time, precise location of the injured animal or carcass, and any other pertinent information. Care should be taken in handling sick or injured specimens to preserve biological materials in the best possible state for later analysis of cause of death, if that occurs. In conjunction with the care of sick or injured endangered or threatened species or preservation of biological materials from a dead animal, the finder has the responsibility to ensure that evidence associated with the specimen is not unnecessarily disturbed. Contact the U.S. Fish and Wildlife Service Law Enforcement Office at (425) 883-8122, or the Service's Washington Fish and Wildlife Office at (360) 753-9440.

CONSERVATION RECOMMENDATIONS

Section 7(a)(1) of the Act directs Federal agencies to utilize their authorities to further the purposes of the Act by carrying out conservation programs for the benefit of endangered and threatened species. Conservation recommendations are discretionary agency activities to minimize or avoid adverse effects of a proposed action on listed species or critical habitat, to help implement recovery plans, or to develop information.

- 1) Explore options for the Corps or other partners to implement aquatic habitat improvement projects that address the sediment from the eroding bluff upstream of action area.

REINITIATION NOTICE

This concludes formal consultation on the action outlined in the request for formal consultation. As provided in 50 CFR 402.16, reinitiation of formal consultation is required where discretionary Federal agency involvement or control over the action has been retained (or is authorized by law) and if: 1) the amount or extent of incidental take is exceeded; 2) new

information reveals effects of the agency action that may affect listed species or critical habitat in a manner or to an extent not considered in this opinion; 3) the agency action is subsequently modified in a manner that causes an effect to the listed species or critical habitat not considered in this opinion; or 4) a new species is listed or critical habitat designated that may be affected by the action. In instances where the amount or extent of incidental take is exceeded, any operations causing such take must cease pending reinitiation.

LITERATURE CITED

- Battin, J., M.W. Wiley, M.H. Ruckelshaus, R.N. Palmer, E. Korb, K.K. Bartz, and H. Imaki. 2007. Projected impacts of climate change on salmon habitat restoration. *Proceedings of the National Academy of Sciences of the United States of America* 104(16):6720-25.
- IPCC (Intergovernmental Panel on Climate Change). 2007a. *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, Pachauri, R.K., and A. Reisinger (eds.)]. IPCC, Geneva, Switzerland. 104pp.
- IPCC (Intergovernmental Panel on Climate Change). 2007b. *Summary for Policymakers*. Pp. 1–18. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, UK, and New York, NY. 996pp.
- IPCC (Intergovernmental Panel on Climate Change). 2011. *Summary for Policymakers*. In: *Intergovernmental Panel on Climate Change Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK, and New York, NY. 29pp.
- IPCC 2014a. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- IPCC 2014b. *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1132 pp.
- ISAB (Independent Scientific Advisory Board). 2007. *Climate change impacts on Columbia River basin fish and wildlife*. ISAB 2007-2. Portland, Oregon.
- Kassler, T.W. and G. Mendel. 2007. *Genetic characterization of Bull Trout from the Walla Walla River Basin*. WDFW, April 2007. 48 pp.

- Mendel, G., J. Trump, and M. Gembala, 2004. Assessment of Salmonids and their Habitat Conditions in the Walla Walla River Basin within Washington, 2003-2004 Annual Report, Project No. 199802000, 137 electronic pages, (BPA Report DOE/BP-00006502-2)
- Mendel, G., C. Fulton, and R. Weldert. An Investigation Into the Migratory Behavior of Bull Trout (*Salvelinus confluentus*) in the Touchet River Basin. WDFW, October 2003. 33 pp.
- Mendel, G., B. Mahoney, R. Weldert, J. Olsen, J. Trump, and A. Fitzgerald. 2014. Walla Walla River Subbasin Salmonid Monitoring and Evaluation Project. 2013 Annual Report for the period 1/1/2013-12/31/2013. BPA Project # 2000-039-00; Report covers work performed under BPA contract #(s) 60695, 61347. 69pp.
- Mendel, G., V. Naef, and D. Karl 1999. Assessment of Salmonid Fishes and Their Habitat Conditions in the Walla Walla River Basin-1998 Annual Report. U.S. Department of Energy, Bonneville Power Administration No. FPA 99-01.
- Mendel, G., D. Karl, and T. Coyle. 2001. Assessment of salmon ids and their habitat conditions in the Walla Walla River Basin: 2000 Annual Report. Project No. 1998-02000, 116 electronic pages, (BPA Report DOEIBP-00000485-1).
- Mendel, G., Trump, J., Gembala, M., Blankenship, S. & Kassler, T. 2007. Assessment of salmonids and their habitat conditions in the Walla Walla River Basin within Washington. p. 181. Portland, OR: Bonneville Power Administration.
- Newcombe, C.P., and D.D. MacDonald. 1991. Effects of suspended sediments on aquatic ecosystems. North American Journal of Fisheries Management 11:72-82.
- Newcombe, C.P. and J.O. Jensen. 1996. Channel Suspended Sediment and Fisheries: a Synthesis for quantitative Assessment of Risk and Impact. N.Am. J. Fish. Management. 16:693-727.
- NMFS (National Marine Fisheries Service). 2009. Middle Columbia River steelhead distinct population segment ESA recovery plan. Northwest Regional Office, Seattle, Washington. [as referenced in BA]
- NMFS (National Marine Fisheries Service). 2011. Endangered Species Act Biological Opinion and Magnuson-Stevens Fishery Conservation and Management Act Essential Fish Habitat Response for the Georgia-Pacific Wauna Mill Transit Dock Repair and Piling Replacement, Columbia River (5th field HUC 1708000307), Clatsop County, Oregon (Corps No.: NWP-2010-587). [as referenced in BA]
- Rieman, B.E., D. Isaak, S. Adams, D. Horan, D. Nagel, C. Luce, and D. Myers. 2007. Anticipated climate warming effects on bull trout habitats and populations across the interior Columbia River Basin. Transactions of the American Fisheries Society 136(6):1552-65.

- Schaller, H., P. Budy, and C. Newlon. Walla Walla River bull trout ten year retrospective Analysis and implications for recovery planning. September 30, 2014. Study funded by U.S. Fish and Wildlife Service and U.S. Geological Survey, Utah Cooperative Fish and Wildlife Research Unit, Department of Watershed Sciences, Utah State University.
- USFWS (U.S. Fish and Wildlife Service). 1998. A framework to assist in making Endangered Species Act determinations of effect for individual or grouped actions at the bull trout subpopulation watershed scale. Western Washington Fish and Wildlife Office, Lacey, WA. 45pp.
- USFWS. 2002. Bull trout (*Salvelinus confluentus*) draft recovery plan. Region 1 U.S. Fish and Wildlife Service Portland, Oregon.
- USFWS. 2004. Bull trout (*Salvelinus confluentus*) draft recovery plan – Chapter 10: Umatilla-Walla Walla. U.S. Fish and Wildlife Service, Portland, Oregon. 148 pp.
- USFWS. 2008. Bull Trout (*Salvelinus confluentus*) 5-Year Review: Summary and Evaluation. 53 pp. Portland, Oregon.
- USFWS. 2008. Bull trout core area status assessment for the Walla Walla River Core Area. September 25, 2008. Upper Columbia Fish and Wildlife Office, Spokane, Washington. 27 pages.
- USFWS 2015a. Recovery plan for the coterminous United States population of bull trout (*Salvelinus confluentus*). Portland, Oregon. xii + 179 pages.
- USFWS 2015b. Bull Trout (*Salvelinus confluentus*) 5-Year Review: Short Form Summary. 8 pp. Boise, Idaho.
- USFWS 2015c. Mid-Columbia Recovery Unit Implementation Plan for Bull Trout (*Salvelinus confluentus*). US Fish and Wildlife Service, Oregon Fish and Wildlife Office, Portland, Oregon. September 2015.
- Wiseman, C.D., M. LeMoine, R. Plotnikoff, J. Diamond, A. Stewart, and S. Cormier. 2010. Identification of Most Probable Stressors to Aquatic Life in the Touchet River, Washington. U.S. Environmental Protection Agency, National Center for Environmental Assessment, Cincinnati, OH. EPA/600/R-08/145.
- World Wildlife Fund (WWF). 2003. Buying time: a user's manual for building resistance and resilience to climate change in natural systems. Editors: L.J. Hansen, J.L Biringer, and J.R.Hoffman.

In Litteris

Driscoll, Diane. 2018. Email from Diane Driscoll regarding communication with John Hook on isolation method and sediment fencing. May 21, 2018.

Hook, John. 2018a. Email response from John Hook, COE, regarding questions and clarifications on project. April 12, 2018.

Hook, John. 2018b. Email response from John Hook, COE, regarding distance of silt fence. May 8, 2018.

Hook, John. 2018c. Email from John Hook, COE, to Diane Driscoll, NMFS, regarding distance of silt fence. May 16, 2018.

APPENDIX A

STATUS OF THE SPECIES: BULL TROUT

Taxonomy

The bull trout (*Salvelinus confluentus*) is a native char found in the coastal and intermountain west of North America. Dolly Varden (*Salvelinus malma*) and bull trout were previously considered a single species and were thought to have coastal and interior forms. However, Cavender (1978, entire) described morphometric, meristic and osteological characteristics of the two forms, and provided evidence of specific distinctions between the two. Despite an overlap in the geographic range of bull trout and Dolly Varden in the Puget Sound area and along the British Columbia coast, there is little evidence of introgression (Haas and McPhail 1991, p. 2191). The Columbia River Basin is considered the region of origin for the bull trout. From the Columbia, dispersal to other drainage systems was accomplished by marine migration and headwater stream capture. Behnke (2002, p. 297) postulated dispersion to drainages east of the continental divide may have occurred through the North and South Saskatchewan Rivers (Hudson Bay drainage) and the Yukon River system. Marine dispersal may have occurred from Puget Sound north to the Fraser, Skeena and Taku Rivers of British Columbia.

Species Description

Bull trout have unusually large heads and mouths for salmonids. Their body colors can vary tremendously depending on their environment, but are often brownish green with lighter (often ranging from pale yellow to crimson) colored spots running along their dorsa and flanks, with spots being absent on the dorsal fin, and light colored to white under bellies. They have white leading edges on their fins, as do other species of char. Bull trout have been measured as large as 103 centimeters (41 inches) in length, with weights as high as 14.5 kilograms (32 pounds) (Fishbase 2015, p. 1). Bull trout may be migratory, moving throughout large river systems, lakes, and even the ocean in coastal populations, or they may be resident, remaining in the same stream their entire lives (Rieman and McIntyre 1993, p. 2; Brenkman and Corbett 2005, p. 1077). Migratory bull trout are typically larger than resident bull trout (USFWS 1998, p. 31668).

Legal Status

The coterminous United States population of the bull trout was listed as threatened on November 1, 1999 (USFWS 1999, entire). The threatened bull trout generally occurs in the Klamath River Basin of south-central Oregon; the Jarbidge River in Nevada; the Willamette River Basin in Oregon; Pacific Coast drainages of Washington, including Puget Sound; major rivers in Idaho, Oregon, Washington, and Montana, within the Columbia River Basin; and the St. Mary-Belly River, east of the Continental Divide in northwestern Montana (Bond 1992, p. 4; Brewin and Brewin 1997, pp. 209-216; Cavender 1978, pp. 165-166; Leary and Allendorf 1997, pp. 715-720).

Throughout its range, the bull trout are threatened by the combined effects of habitat degradation, fragmentation, and alterations associated with dewatering, road construction and

maintenance, mining, grazing, the blockage of migratory corridors by dams or other diversion structures, poor water quality, entrainment (a process by which aquatic organisms are pulled through a diversion or other device) into diversion channels, and introduced non-native species (USFWS 1999, p. 58910). Although all salmonids are likely to be affected by climate change, bull trout are especially vulnerable given that spawning and rearing are constrained by their location in upper watersheds and the requirement for cold water temperatures (Battin et al. 2007, entire; Rieman et al. 2007, entire; Porter and Nelitz. 2009, pages 4-8). Poaching and incidental mortality of bull trout during other targeted fisheries are additional threats.

Life History

The iteroparous reproductive strategy of bull trout has important repercussions for the management of this species. Bull trout require passage both upstream and downstream, not only for repeat spawning but also for foraging. Most fish ladders, however, were designed specifically for anadromous semelparous salmonids (fishes that spawn once and then die, and require only one-way passage upstream). Therefore, even dams or other barriers with fish passage facilities may be a factor in isolating bull trout populations if they do not provide a downstream passage route. Additionally, in some core areas, bull trout that migrate to marine waters must pass both upstream and downstream through areas with net fisheries at river mouths. This can increase the likelihood of mortality to bull trout during these spawning and foraging migrations.

Growth varies depending upon life-history strategy. Resident adults range from 6 to 12 inches total length, and migratory adults commonly reach 24 inches or more (Goetz 1989, p. 30; Pratt 1985, pp. 28-34). The largest verified bull trout is a 32-pound specimen caught in Lake Pend Oreille, Idaho, in 1949 (Simpson and Wallace 1982, p. 95).

Bull trout typically spawn from August through November during periods of increasing flows and decreasing water temperatures. Preferred spawning habitat consists of low-gradient stream reaches with loose, clean gravel (Fraley and Shepard 1989, p. 141). Redds are often constructed in stream reaches fed by springs or near other sources of cold groundwater (Goetz 1989, pp. 15-16; Pratt 1992, pp. 6-7; Rieman and McIntyre 1996, p. 133). Depending on water temperature, incubation is normally 100 to 145 days (Pratt 1992, p. 1). After hatching, fry remain in the substrate, and time from egg deposition to emergence may surpass 200 days. Fry normally emerge from early April through May, depending on water temperatures and increasing stream flows (Pratt 1992, p. 1; Ratliff and Howell 1992, p. 10).

Early life stages of fish, specifically the developing embryo, require the highest inter-gravel dissolved oxygen (IGDO) levels, and are the most sensitive life stage to reduced oxygen levels. The oxygen demand of embryos depends on temperature and on stage of development, with the greatest IGDO required just prior to hatching.

A literature review conducted by the Washington Department of Ecology (WDOE 2002, p. 9) indicates that adverse effects of lower oxygen concentrations on embryo survival are magnified as temperatures increase above optimal (for incubation). Normal oxygen levels seen in rivers used by bull trout during spawning ranged from 8 to 12 mg/L (in the gravel), with corresponding

instream levels of 10 to 11.5 mg/L (Stewart et al. 2007, p. 10). In addition, IGDO concentrations, water velocities in the water column, and especially the intergravel flow rate, are interrelated variables that affect the survival of incubating embryos (ODEQ 1995, Ch 2 pp. 23-24). Due to a long incubation period of 220+ days, bull trout are particularly sensitive to adequate IGDO levels. An IGDO level below 8 mg/L is likely to result in mortality of eggs, embryos, and fry.

Population Dynamics

Population Structure

Bull trout exhibit both resident and migratory life history strategies. Both resident and migratory forms may be found together, and either form may produce offspring exhibiting either resident or migratory behavior (Rieman and McIntyre 1993, p. 2). Resident bull trout complete their entire life cycle in the tributary (or nearby) streams in which they spawn and rear. The resident form tends to be smaller than the migratory form at maturity and also produces fewer eggs (Goetz 1989, p. 15). Migratory bull trout spawn in tributary streams where juvenile fish rear 1 to 4 years before migrating to either a lake (adfluvial form), river (fluvial form) (Fraley and Shepard 1989, p. 138; Goetz 1989, p. 24), or saltwater (anadromous form) to rear as subadults and to live as adults (Brenkman and Corbett 2005, entire; McPhail and Baxter 1996, p. i; WDFW et al. 1997, p. 16). Bull trout normally reach sexual maturity in 4 to 7 years and may live longer than 12 years. They are iteroparous (they spawn more than once in a lifetime). Repeat- and alternate-year spawning has been reported, although repeat-spawning frequency and post-spawning mortality are not well documented (Fraley and Shepard 1989, p. 135; Leathe and Graham 1982, p. 95; Pratt 1992, p. 8; Rieman and McIntyre 1996, p. 133).

Bull trout are naturally migratory, which allows them to capitalize on temporally abundant food resources and larger downstream habitats. Resident forms may develop where barriers (either natural or manmade) occur or where foraging, migrating, or overwintering habitats for migratory fish are minimized (Brenkman and Corbett 2005, pp. 1075-1076; Goetz et al. 2004, p. 105). For example, multiple life history forms (e.g., resident and fluvial) and multiple migration patterns have been noted in the Grande Ronde River (Baxter 2002, pp. 96, 98-106). Parts of this river system have retained habitat conditions that allow free movement between spawning and rearing areas and the mainstem Snake River. Such multiple life history strategies help to maintain the stability and persistence of bull trout populations to environmental changes. Benefits to migratory bull trout include greater growth in the more productive waters of larger streams, lakes, and marine waters; greater fecundity resulting in increased reproductive potential; and dispersing the population across space and time so that spawning streams may be recolonized should local populations suffer a catastrophic loss (Frissell 1999, pp. 861-863; MBTSG 1998, p. 13; Rieman and McIntyre 1993, pp. 2-3). In the absence of the migratory bull trout life form, isolated populations cannot be replenished when disturbances make local habitats temporarily unsuitable. Therefore, the range of the species is diminished, and the potential for a greater reproductive contribution from larger size fish with higher fecundity is lost (Rieman and McIntyre 1993, p. 2).

Whitesel et al. (2004, p. 2) noted that although there are multiple resources that contribute to the subject, Spruell et al. (2003, entire) best summarized genetic information on bull trout population

structure. Spruell et al. (2003, entire) analyzed 1,847 bull trout from 65 sampling locations, four located in three coastal drainages (Klamath, Queets, and Skagit Rivers), one in the Saskatchewan River drainage (Belly River), and 60 scattered throughout the Columbia River Basin. They concluded that there is a consistent pattern among genetic studies of bull trout, regardless of whether examining allozymes, mitochondrial DNA, or most recently microsatellite loci. Typically, the genetic pattern shows relatively little genetic variation within populations, but substantial divergence among populations. Microsatellite loci analysis supports the existence of at least three major genetically differentiated groups (or evolutionary lineages) of bull trout (Spruell et al. 2003, p. 17). They were characterized as:

- i. "Coastal", including the Deschutes River and all of the Columbia River drainage downstream, as well as most coastal streams in Washington, Oregon, and British Columbia. A compelling case also exists that the Klamath Basin represents a unique evolutionary lineage within the coastal group.
- ii. "Snake River", which also included the John Day, Umatilla, and Walla Walla rivers. Despite close proximity of the John Day and Deschutes Rivers, a striking level of divergence between bull trout in these two systems was observed.
- iii. "Upper Columbia River" which includes the entire basin in Montana and northern Idaho. A tentative assignment was made by Spruell et al. (2003, p. 25) of the Saskatchewan River drainage populations (east of the continental divide), grouping them with the upper Columbia River group.

Spruell et al. (2003, p. 17) noted that within the major assemblages, populations were further subdivided, primarily at the level of major river basins. Taylor et al. (1999, entire) surveyed bull trout populations, primarily from Canada, and found a major divergence between inland and coastal populations. Costello et al. (2003, p. 328) suggested the patterns reflected the existence of two glacial refugia, consistent with the conclusions of Spruell et al. (2003, p. 26) and the biogeographic analysis of Haas and McPhail (2001, entire). Both Taylor et al. (1999, p. 1166) and Spruell et al. (2003, p. 21) concluded that the Deschutes River represented the most upstream limit of the coastal lineage in the Columbia River Basin.

More recently, the Service identified additional genetic units within the coastal and interior lineages (Ardren et al. 2011, p. 18). Based on a recommendation in the Service's 5-year review of the species' status (USFWS 2008a, p. 45), the Service reanalyzed the 27 recovery units identified in the draft bull trout recovery plan (USFWS 2002a, p. 48) by utilizing, in part, information from previous genetic studies and new information from additional analysis (Ardren et al. 2011, entire). In this examination, the Service applied relevant factors from the joint Service and National Marine Fisheries Service Distinct Population Segment (DPS) policy (USFWS 1996, entire) and subsequently identified six draft recovery units that contain assemblages of core areas that retain genetic and ecological integrity across the range of bull trout in the coterminous United States. These six draft recovery units were used to inform designation of critical habitat for bull trout by providing a context for deciding what habitats are essential for recovery (USFWS 2010, p. 63898). The six draft recovery units identified for bull trout in the coterminous United States include: Coastal, Klamath, Mid-Columbia, Columbia Headwaters, Saint Mary, and Upper Snake. These six draft recovery units were also identified in the Service's revised recovery plan (USFWS 2015, p. vii) and designated as final recovery units.

Population Dynamics

Although bull trout are widely distributed over a large geographic area, they exhibit a patchy distribution, even in pristine habitats (Rieman and McIntyre 1993, p. 4). Increased habitat fragmentation reduces the amount of available habitat and increases isolation from other populations of the same species (Saunders et al. 1991, entire). Burkey (1989, entire) concluded that when species are isolated by fragmented habitats, low rates of population growth are typical in local populations and their probability of extinction is directly related to the degree of isolation and fragmentation. Without sufficient immigration, growth for local populations may be low and probability of extinction high (Burkey 1989, entire; Burkey 1995, entire). Metapopulation concepts of conservation biology theory have been suggested relative to the distribution and characteristics of bull trout, although empirical evidence is relatively scant (Rieman and McIntyre 1993, p. 15; Dunham and Rieman 1999, entire; Rieman and Dunham 2000, entire). A metapopulation is an interacting network of local populations with varying frequencies of migration and gene flow among them (Meffe and Carroll 1994, pp. 189-190). For inland bull trout, metapopulation theory is likely most applicable at the watershed scale where habitat consists of discrete patches or collections of habitat capable of supporting local populations; local populations are for the most part independent and represent discrete reproductive units; and long-term, low-rate dispersal patterns among component populations influences the persistence of at least some of the local populations (Rieman and Dunham 2000, entire). Ideally, multiple local populations distributed throughout a watershed provide a mechanism for spreading risk because the simultaneous loss of all local populations is unlikely. However, habitat alteration, primarily through the construction of impoundments, dams, and water diversions has fragmented habitats, eliminated migratory corridors, and in many cases isolated bull trout in the headwaters of tributaries (Rieman and Clayton 1997, pp. 10-12; Dunham and Rieman 1999, p. 645; Spruell et al. 1999, pp. 118-120; Rieman and Dunham 2000, p. 55).

Human-induced factors as well as natural factors affecting bull trout distribution have likely limited the expression of the metapopulation concept for bull trout to patches of habitat within the overall distribution of the species (Dunham and Rieman 1999, entire). However, despite the theoretical fit, the relatively recent and brief time period during which bull trout investigations have taken place does not provide certainty as to whether a metapopulation dynamic is occurring (e.g., a balance between local extirpations and recolonizations) across the range of the bull trout or whether the persistence of bull trout in large or closely interconnected habitat patches (Dunham and Rieman 1999, entire) is simply reflective of a general deterministic trend towards extinction of the species where the larger or interconnected patches are relics of historically wider distribution (Rieman and Dunham 2000, pp. 56-57). Recent research (Whiteley et al. 2003, entire) does, however, provide genetic evidence for the presence of a metapopulation process for bull trout, at least in the Boise River Basin of Idaho.

Habitat Characteristics

Bull trout have more specific habitat requirements than most other salmonids (Rieman and McIntyre 1993, p. 4). Habitat components that influence bull trout distribution and abundance include water temperature, cover, channel form and stability, valley form, spawning and rearing substrate, and migratory corridors (Fraleley and Shepard 1989, entire; Goetz 1989, pp. 23, 25; Hoelscher and Bjornn 1989, pp. 19, 25; Howell and Buchanan 1992, pp. 30, 32; Pratt 1992,

entire; Rich 1996, p. 17; Rieman and McIntyre 1993, pp. 4-6; Rieman and McIntyre 1995, entire; Sedell and Everest 1991, entire; Watson and Hillman 1997, entire). Watson and Hillman (1997, pp. 247-250) concluded that watersheds must have specific physical characteristics to provide the habitat requirements necessary for bull trout to successfully spawn and rear and that these specific characteristics are not necessarily present throughout these watersheds. Because bull trout exhibit a patchy distribution, even in pristine habitats (Rieman and McIntyre 1993, pp. 4-6), bull trout should not be expected to simultaneously occupy all available habitats.

Migratory corridors link seasonal habitats for all bull trout life histories. The ability to migrate is important to the persistence of bull trout (Rieman and McIntyre 1993, p. 2). Migrations facilitate gene flow among local populations when individuals from different local populations interbreed or stray to nonnatal streams. Local populations that are extirpated by catastrophic events may also become reestablished by bull trout migrants. However, it is important to note that the genetic structuring of bull trout indicates there is limited gene flow among bull trout populations, which may encourage local adaptation within individual populations, and that reestablishment of extirpated populations may take a long time (Rieman and McIntyre 1993, p. 2; Spruell et al. 1999, entire). Migration also allows bull trout to access more abundant or larger prey, which facilitates growth and reproduction. Additional benefits of migration and its relationship to foraging are discussed below under "Diet."

Cold water temperatures play an important role in determining bull trout habitat quality, as these fish are primarily found in colder streams, and spawning habitats are generally characterized by temperatures that drop below 9 °C in the fall (Fraley and Shepard 1989, p. 137; Pratt 1992, p. 5; Rieman and McIntyre 1993, p. 2).

Thermal requirements for bull trout appear to differ at different life stages. Spawning areas are often associated with cold-water springs, groundwater infiltration, and the coldest streams in a given watershed (Pratt 1992, pp 7-8; Rieman and McIntyre 1993, p. 7). Optimum incubation temperatures for bull trout eggs range from 2 °C to 6 °C whereas optimum water temperatures for rearing range from about 6 °C to 10 °C (Buchanan and Gregory 1997, p. 4; Goetz 1989, p. 22). In Granite Creek, Idaho, Bonneau and Scarnecchia (1996, entire) observed that juvenile bull trout selected the coldest water available in a plunge pool, 8 °C to 9 °C, within a temperature gradient of 8 °C to 15 °C. In a landscape study relating bull trout distribution to maximum water temperatures, Dunham et al. (2003, p. 900) found that the probability of juvenile bull trout occurrence does not become high (i.e., greater than 0.75) until maximum temperatures decline to 11 °C to 12 °C.

Although bull trout are found primarily in cold streams, occasionally these fish are found in larger, warmer river systems throughout the Columbia River basin (Buchanan and Gregory 1997, p. 2; Fraley and Shepard 1989, pp. 133, 135; Rieman and McIntyre 1993, pp. 3-4; Rieman and McIntyre 1995, p. 287). Availability and proximity of cold water patches and food productivity can influence bull trout ability to survive in warmer rivers (Myrick 2002, pp. 6 and 13).

All life history stages of bull trout are associated with complex forms of cover, including large woody debris, undercut banks, boulders, and pools (Fraley and Shepard 1989, p. 137; Goetz 1989, p. 19; Hoelscher and Bjornn 1989, p. 38; Pratt 1992, entire; Rich 1996, pp. 4-5; Sedell and Everest 1991, entire; Sexauer and James 1997, entire; Thomas 1992, pp. 4-6; Watson and Hillman 1997, p. 238). Maintaining bull trout habitat requires natural stability of stream channels and maintenance of natural flow patterns (Rieman and McIntyre 1993, pp. 5-6). Juvenile and adult bull trout frequently inhabit side channels, stream margins, and pools with suitable cover (Sexauer and James 1997, p. 364). These areas are sensitive to activities that directly or indirectly affect stream channel stability and alter natural flow patterns. For example, altered stream flow in the fall may disrupt bull trout during the spawning period, and channel instability may decrease survival of eggs and young juveniles in the gravel from winter through spring (Fraley and Shepard 1989, p. 141; Pratt 1992, p. 6; Pratt and Huston 1993, p. 70). Pratt (1992, p. 6) indicated that increases in fine sediment reduce egg survival and emergence.

Diet

Bull trout are opportunistic feeders, with food habits primarily a function of size and life-history strategy. Fish growth depends on the quantity and quality of food that is eaten, and as fish grow their foraging strategy changes as their food changes, in quantity, size, or other characteristics (Quinn 2005, pp. 195-200). Resident and juvenile migratory bull trout prey on terrestrial and aquatic insects, macrozooplankton, and small fish (Boag 1987, p. 58; Donald and Alger 1993, pp. 242-243; Goetz 1989, pp. 33-34). Subadult and adult migratory bull trout feed on various fish species (Donald and Alger 1993, pp. 241-243; Fraley and Shepard 1989, pp. 135, 138; Leathe and Graham 1982, pp. 13, 50-56). Bull trout of all sizes other than fry have been found to eat fish half their length (Beauchamp and VanTassell 2001, p. 204). In nearshore marine areas of western Washington, bull trout feed on Pacific herring (*Clupea pallasii*), Pacific sand lance (*Ammodytes hexapterus*), and surf smelt (*Hypomesus pretiosus*) (Goetz et al. 2004, p. 105; WDFW et al. 1997, p. 23).

Bull trout migration and life history strategies are closely related to their feeding and foraging strategies. Migration allows bull trout to access optimal foraging areas and exploit a wider variety of prey resources. For example, in the Skagit River system, anadromous bull trout make migrations as long as 121 miles between marine foraging areas in Puget Sound and headwater spawning grounds, foraging on salmon eggs and juvenile salmon along their migration route (WDFW et al. 1997, p. 25). Anadromous bull trout also use marine waters as migration corridors to reach seasonal habitats in non-natal watersheds to forage and possibly overwinter (Brenkman and Corbett 2005, pp. 1078-1079; Goetz et al. 2004, entire).

Status and Distribution

Distribution and Demography

The historical range of bull trout includes major river basins in the Pacific Northwest at about 41 to 60 degrees North latitude, from the southern limits in the McCloud River in northern California and the Jarbidge River in Nevada to the headwaters of the Yukon River in the Northwest Territories, Canada (Cavender 1978, pp. 165-166; Bond 1992, p. 2). To the west, the bull trout's range includes Puget Sound, various coastal rivers of British Columbia, Canada, and southeast Alaska (Bond 1992, p. 2). Bull trout occur in portions of the Columbia River and

tributaries within the basin, including its headwaters in Montana and Canada. Bull trout also occur in the Klamath River basin of south-central Oregon. East of the Continental Divide, bull trout are found in the headwaters of the Saskatchewan River in Alberta and Montana and in the MacKenzie River system in Alberta and British Columbia, Canada (Cavender 1978, pp. 165-166; Brewin et al. 1997, entire).

Each of the following recovery units (below) is necessary to maintain the bull trout's distribution, as well as its genetic and phenotypic diversity, all of which are important to ensure the species' resilience to changing environmental conditions. No new local populations have been identified and no local populations have been lost since listing.

Coastal Recovery Unit

The Coastal Recovery Unit is located within western Oregon and Washington. Major geographic regions include the Olympic Peninsula, Puget Sound, and Lower Columbia River basins. The Olympic Peninsula and Puget Sound geographic regions also include their associated marine waters (Puget Sound, Hood Canal, Strait of Juan de Fuca, and Pacific Coast), which are critical in supporting the anadromous¹ life history form, unique to the Coastal Recovery Unit. The Coastal Recovery Unit is also the only unit that overlaps with the distribution of Dolly Varden (*Salvelinus malma*) (Ardren et al. 2011), another native char species that looks very similar to the bull trout (Haas and McPhail 1991). The two species have likely had some level of historic introgression in this part of their range (Redenbach and Taylor 2002). The Lower Columbia River major geographic region includes the lower mainstem Columbia River, an important migratory waterway essential for providing habitat and population connectivity within this region. In the Coastal Recovery Unit, there are 21 existing bull trout core areas which have been designated, including the recently reintroduced Clackamas River population, and 4 core areas have been identified that could be re-established. Core areas within the recovery unit are distributed among these three major geographic regions (Puget Sound also includes one core area that is actually part of the lower Fraser River system in British Columbia, Canada) (USFWS 2015a, p. A-1).

The current demographic status of bull trout in the Coastal Recovery Unit is variable across the unit. Populations in the Puget Sound region generally tend to have better demographic status, followed by the Olympic Peninsula, and finally the Lower Columbia River region. However, population strongholds do exist across the three regions. The Lower Skagit River and Upper Skagit River core areas in the Puget Sound region likely contain two of the most abundant bull trout populations with some of the most intact habitat within this recovery unit. The Lower Deschutes River core area in the Lower Columbia River region also contains a very abundant bull trout population and has been used as a donor stock for re-establishing the Clackamas River population (USFWS 2015a, p. A-6).

Puget Sound Region

In the Puget Sound region, bull trout populations are concentrated along the eastern side of Puget Sound with most core areas concentrated in central and northern Puget Sound. Although the Chilliwack River core area is considered part of this region, it is technically connected to the Fraser River system and is transboundary with British Columbia making its

¹ Anadromous: Life history pattern of spawning and rearing in fresh water and migrating to salt water areas to mature.

distribution unique within the region. Most core areas support a mix of anadromous and fluvial life history forms, with at least two core areas containing a natural adfluvial life history (Chilliwack River core area [Chilliwack Lake] and Chester Morse Lake core area). Overall demographic status of core areas generally improves as you move from south Puget Sound to north Puget Sound. Although comprehensive trend data are lacking, the current condition of core areas within this region are likely stable overall, although some at depressed abundances. Two core areas (Puyallup River and Stillaguamish River) contain local populations at either very low abundances (Upper Puyallup and Mowich Rivers) or that have likely become locally extirpated (Upper Deer Creek, South Fork Canyon Creek, and Greenwater River). Connectivity among and within core areas of this region is generally intact. Most core areas in this region still have significant amounts of headwater habitat within protected and relatively pristine areas (e.g., North Cascades National Park, Mount Rainier National Park, Skagit Valley Provincial Park, Manning Provincial Park, and various wilderness or recreation areas) (USFWS 2015a, p. A-7).

Olympic Peninsula Region

In the Olympic Peninsula region, distribution of core areas is somewhat disjunct, with only one located on the west side of Hood Canal on the eastern side of the peninsula, two along the Strait of Juan de Fuca on the northern side of the peninsula, and three along the Pacific Coast on the western side of the peninsula. Most core areas support a mix of anadromous and fluvial life history forms, with at least one core area also supporting a natural adfluvial life history (Quinault River core area [Quinault Lake]). Demographic status of core areas is poorest in Hood Canal and Strait of Juan de Fuca, while core areas along the Pacific Coast of Washington likely have the best demographic status in this region. The connectivity between core areas in these disjunct regions is believed to be naturally low due to the geographic distance between them.

Internal connectivity is currently poor within the Skokomish River core area (Hood Canal) and is being restored in the Elwha River core area (Strait of Juan de Fuca). Most core areas in this region still have their headwater habitats within relatively protected areas (Olympic National Park and wilderness areas) (USFWS 2015a, p. A-7).

Lower Columbia River Region

In the Lower Columbia River region, the majority of core areas are distributed along the Cascade Crest on the Oregon side of the Columbia River. Only two of the seven core areas in this region are in Washington. Most core areas in the region historically supported a fluvial life history form, but many are now adfluvial due to reservoir construction. However, there is at least one core area supporting a natural adfluvial life history (Odell Lake) and one supporting a natural, isolated, resident life history (Klickitat River [West Fork Klickitat]). Status is highly variable across this region, with one relative stronghold (Lower Deschutes core area) existing on the Oregon side of the Columbia River. The Lower Columbia River region also contains three watersheds (North Santiam River, Upper Deschutes River, and White Salmon River) that could potentially become re-established core areas within the Coastal Recovery Unit. Although the South Santiam River has been identified as a historic core area, there remains uncertainty as to whether or not historical observations of bull trout represented a self-sustaining population. Current habitat conditions in the South Santiam River are thought to be unable to support bull

trout spawning and rearing. Adult abundances within the majority of core areas in this region are relatively low, generally 300 or fewer individuals.

Most core populations in this region are not only isolated from one another due to dams or natural barriers, but they are internally fragmented as a result of manmade barriers. Local populations are often disconnected from one another or from potential foraging habitat. In the Coastal Recovery Unit, adult abundance may be lowest in the Hood River and Odell Lake core areas, which each contain fewer than 100 adults. Bull trout were reintroduced in the Middle Fork Willamette River in 1990 above Hills Creek Reservoir. Successful reproduction was first documented in 2006, and has occurred each year since (USFWS 2015a, p. A-8). Natural reproducing populations of bull trout are present in the McKenzie River basin (USFWS 2008d, pp. 65-67). Bull trout were more recently reintroduced into the Clackamas River basin in the summer of 2011 after an extensive feasibility analysis (Shively et al. 2007, Hudson et al. 2015). Bull trout from the Lower Deschutes core area are being utilized for this reintroduction effort (USFWS 2015a, p. A-8).

Klamath Recovery Unit

Bull trout in the Klamath Recovery Unit have been isolated from other bull trout populations for the past 10,000 years and are recognized as evolutionarily and genetically distinct (Minckley et al. 1986; Leary et al. 1993; Whitesel et al. 2004; USFWS 2008a; Ardren et al. 2011). As such, there is no opportunity for bull trout in another recovery unit to naturally re-colonize the Klamath Recovery Unit if it were to become extirpated. The Klamath Recovery Unit lies at the southern edge of the species range and occurs in an arid portion of the range of bull trout. Bull trout were once widespread within the Klamath River basin (Gilbert 1897; Dambacher et al. 1992; Ziller 1992; USFWS 2002b), but habitat degradation and fragmentation, past and present land use practices, agricultural water diversions, and past fisheries management practices have greatly reduced their distribution. Bull trout abundance also has been severely reduced, and the remaining populations are highly fragmented and vulnerable to natural or manmade factors that place them at a high risk of extirpation (USFWS 2002b). The presence of nonnative brook trout (*Salvelinus fontinalis*), which compete and hybridize with bull trout, is a particular threat to bull trout persistence throughout the Klamath Recovery Unit (USFWS 2015b, pp. B-3-4).

Upper Klamath Lake Core Area

The Upper Klamath Lake core area comprises two bull trout local populations (Sun Creek and Threemile Creek). These local populations likely face an increased risk of extirpation because they are isolated and not interconnected with each other. Extirpation of other local populations in the Upper Klamath Lake core area has occurred in recent times (1970s). Populations in this core area are genetically distinct from those in the other two core areas in the Klamath Recovery Unit (USFWS 2008b), and in comparison, genetic variation within this core area is lowest. The two local populations have been isolated by habitat fragmentation and have experienced population bottlenecks. As such, currently unoccupied habitat is needed to restore connectivity between the two local populations and to establish additional populations. This unoccupied habitat includes canals, which now provide the only means of connectivity as migratory corridors. Providing full volitional connectivity for bull trout, however, also introduces the risk of invasion by brook trout, which are abundant in this core area.

Bull trout in the Upper Klamath Lake core area formerly occupied Annie Creek, Sevenmile Creek, Cherry Creek, and Fort Creek, but are now extirpated from these locations. The last remaining local populations, Sun Creek and Threemile Creek, have received focused attention. Brook trout have been removed from bull trout occupied reaches, and these reaches have been intentionally isolated to prevent brook trout reinvasion. As such, over the past few generations these populations have become stable and have increased in distribution and abundance. In 1996, the Threemile Creek population had approximately 50 fish that occupied a 1.4-km (0.9-mile) reach (USFWS 2002b). In 2012, a mark-resight population estimate was completed in Threemile Creek, which indicated an abundance of 577 (95 percent confidence interval = 475 to 679) age-1+ fish (ODFW 2012). In addition, the length of the distribution of bull trout in Threemile Creek had increased to 2.7 km (1.7 miles) by 2012 (USFWS unpublished data). Between 1989 and 2010, bull trout abundance in Sun Creek increased approximately tenfold (from approximately 133 to 1,606 age-1+ fish) and distribution increased from approximately 1.9 km (1.2 miles) to 11.2 km (7.0 miles) (Buktenica et al. 2013) (USFWS 2015b, p. B-5).

Sycan River Core Area

The Sycan River core area is comprised of one local population, Long Creek. Long Creek likely faces greater risk of extirpation because it is the only remaining local population due to extirpation of all other historic local populations. Bull trout previously occupied Calahan Creek, Coyote Creek, and the Sycan River, but are now extirpated from these locations (Light et al. 1996). This core area's local population is genetically distinct from those in the other two core areas (USFWS 2008b). This core area also is essential for recovery because bull trout in this core area exhibit both resident² and fluvial life histories, which are important for representing diverse life history expression in the Klamath Recovery Unit. Migratory bull trout are able to grow larger than their resident counterparts, resulting in greater fecundity and higher reproductive potential (Rieman and McIntyre 1993). Migratory life history forms also have been shown to be important for population persistence and resilience (Dunham et al. 2008). The last remaining population (Long Creek) has received focused attention in an effort to ensure it is not also extirpated. In 2006, two weirs were removed from Long Creek, which increased the amount of occupied FMO habitat by 3.2 km (2.0 miles). Bull trout currently occupy approximately 3.5 km (2.2 miles) of spawning/rearing habitat, including a portion of an unnamed tributary to upper Long Creek, and seasonally use 25.9 km (16.1 miles) of FMO habitat. Brook trout also inhabit Long Creek and have been the focus of periodic removal efforts. No recent statistically rigorous population estimate has been completed for Long Creek; however, the 2002 Draft Bull Trout Recovery Plan reported a population estimate of 842 individuals (USFWS 2002b). Currently unoccupied habitat is needed to establish additional local populations, although brook trout are widespread in this core area and their management will need to be considered in future recovery efforts. In 2014, the Klamath Falls Fish and Wildlife Office of the Service established an agreement with the U.S. Geological Survey to undertake a structured decision making process to assist with recovery planning of bull trout populations in the Sycan River core area (USFWS 2015b, p. B-6).

Upper Sprague River Core Area

The Upper Sprague River core area comprises five bull trout local populations, placing the core area at an intermediate risk of extinction. The five local populations include Boulder Creek,

² Resident: Life history pattern of residing in tributary streams for the fish's entire life without migrating.

Dixon Creek, Deming Creek, Leonard Creek, and Brownsworth Creek. These local populations may face a higher risk of extirpation because not all are interconnected. Bull trout local populations in this core area are genetically distinct from those in the other two Klamath Recovery Unit core areas (USFWS 2008b). Migratory bull trout have occasionally been observed in the North Fork Sprague River (USFWS 2002b). Therefore, this core area also is essential for recovery in that bull trout here exhibit a resident life history and likely a fluvial life history, which are important for conserving diverse life history expression in the Klamath Recovery Unit as discussed above for the Sycan River core area.

The Upper Sprague River core area population of bull trout has experienced a decline from historic levels, although less is known about historic occupancy in this core area. Bull trout are reported to have historically occupied the South Fork Sprague River, but are now extirpated from this location (Buchanan et al. 1997). The remaining five populations have received focused attention. Although brown trout (*Salmo trutta*) co-occur with bull trout and exist in adjacent habitats, brook trout do not overlap with existing bull trout populations. Efforts have been made to increase connectivity of existing bull trout populations by replacing culverts that create barriers. Thus, over the past few generations, these populations have likely been stable and increased in distribution. Population abundance has been estimated recently for Boulder Creek (372 + 62 percent; Hartill and Jacobs 2007), Dixon Creek (20 + 60 percent; Hartill and Jacobs 2007), Deming Creek (1,316 + 342; Moore 2006), and Leonard Creek (363 + 37 percent; Hartill and Jacobs 2007). No statistically rigorous population estimate has been completed for the Brownsworth Creek local population; however, the 2002 Draft Bull Trout Recovery Plan reported a population estimate of 964 individuals (USFWS 2002b). Additional local populations need to be established in currently unoccupied habitat within the Upper Sprague River core area, although brook trout are widespread in this core area and will need to be considered in future recovery efforts (USFWS 2015b, p. B-7).

Mid-Columbia Recovery Unit

The Mid-Columbia Recovery Unit (RU) comprises 24 bull trout core areas, as well as 2 historically occupied core areas and 1 research needs area. The Mid-Columbia RU is recognized as an area where bull trout have co-evolved with salmon, steelhead, lamprey, and other fish populations. Reduced fish numbers due to historic overfishing and land management changes have caused changes in nutrient abundance for resident migratory fish like the bull trout. The recovery unit is located within eastern Washington, eastern Oregon, and portions of central Idaho. Major drainages include the Methow River, Wenatchee River, Yakima River, John Day River, Umatilla River, Walla Walla River, Grande Ronde River, Imnaha River, Clearwater River, and smaller drainages along the Snake River and Columbia River (USFWS 2015c, p. C-1).

The Mid-Columbia RU can be divided into four geographic regions the Lower Mid-Columbia, which includes all core areas that flow into the Columbia River below its confluence with the 1) Snake River; 2) the Upper Mid-Columbia, which includes all core areas that flow into the Columbia River above its confluence with the Snake River; 3) the Lower Snake, which includes all core areas that flow into the Snake River between its confluence with the Columbia River and Hells Canyon Dam; and 4) the Mid-Snake, which includes all core areas in the Mid-Columbia RU that flow into the Snake River above Hells Canyon Dam. These geographic regions are

composed of neighboring core areas that share similar bull trout genetic, geographic (hydrographic), and/or habitat characteristics. Conserving bull trout in geographic regions allows for the maintenance of broad representation of genetic diversity, provides neighboring core areas with potential source populations in the event of local extirpations, and provides a broad array of options among neighboring core areas to contribute recovery under uncertain environmental change (USFWS 2015c, pp. C-1-2).

The current demographic status of bull trout in the Mid-Columbia Recovery Unit is highly variable at both the RU and geographic region scale. Some core areas, such as the Umatilla, Asotin, and Powder Rivers, contain populations so depressed they are likely suffering from the deleterious effects of small population size. Conversely, strongholds do exist within the recovery unit, predominantly in the Lower Snake geographic area. Populations in the Imnaha, Little Minam, Clearwater, and Wenaha Rivers are likely some of the most abundant. These populations are all completely or partially within the bounds of protected wilderness areas and have some of the most intact habitat in the recovery unit. Status in some core areas is relatively unknown, but all indications in these core areas suggest population trends are declining, particularly in the core areas of the John Day Basin (USFWS 2015c, p. C-5).

Lower Mid-Columbia Region

In the Lower Mid-Columbia Region, core areas are distributed along the western portion of the Blue Mountains in Oregon and Washington. Only one of the six core areas is located completely in Washington. Demographic status is highly variable throughout the region. Status is the poorest in the Umatilla and Middle Fork John Day Core Areas. However, the Walla Walla River core area contains nearly pristine habitats in the headwater spawning areas and supports the most abundant populations in the region. Most core areas support both a resident and fluvial life history; however, recent evidence suggests a significant decline in the resident and fluvial life history in the Umatilla River and John Day core areas respectively. Connectivity between the core areas of the Lower Mid-Columbia Region is unlikely given conditions in the connecting FMO habitats. Connection between the Umatilla, Walla Walla and Touchet core areas is uncommon but has been documented, and connectivity is possible between core areas in the John Day Basin. Connectivity between the John Day core areas and Umatilla/Walla Walla/Touchet core areas is unlikely (USFWS 2015c, pp. C-5-6).

Upper Mid-Columbia Region

In the Upper Mid-Columbia Region, core areas are distributed along the eastern side of the Cascade Mountains in Central Washington and part of northeastern Washington. This area contains five core areas (Yakima, Wenatchee, Entiat, Methow, and South Salmo), the Lake Chelan historic core area, Northeast Washington Research Needs Area, and the Chelan River, Okanogan River, and Columbia River FMO areas. The core area populations are generally considered migratory, though they currently express both migratory (fluvial and adfluvial) and resident forms. Residents are located both above and below natural barriers (*i.e.*, Early Winters Creek above a natural falls; and Ahtanum in the Yakima likely due to long lack of connectivity from irrigation withdrawal). In terms of uniqueness and connectivity, the genetics baseline, radio-telemetry, and PIT tag studies identified unique local populations in all core areas. Movement patterns within the core areas; between the lower river, lakes, and other core areas; and between the Chelan, Okanogan, and Columbia River FMO occurs regularly for some of the

Wenatchee, Entiat, and Methow core area populations. This type of connectivity has been displayed by one or more fish, typically in non-spawning movements within FMO. More recently, connectivity has been observed between the Entiat and Yakima core areas by a juvenile bull trout tagged in the Entiat moving in to the Yakima at Prosser Dam and returning at an adult size back to the Entiat. Genetics baselines identify unique populations in all four core areas (USFWS 2015c, p. C-6).

The demographic status is variable in the Upper-Mid Columbia region and ranges from good to very poor. The Service's 2008 5-year Review and Conservation Status Assessment described the Methow and Yakima Rivers at risk, with a rapidly declining trend. The Entiat River was listed at risk with a stable trend, and the Wenatchee River as having a potential risk, and with a stable trend. Currently, the Entiat River is considered to be declining rapidly due to much reduced redd counts. The Wenatchee River is able to exhibit all freshwater life histories with connectivity to Lake Wenatchee, the Wenatchee River and all its local populations, and to the Columbia River and/or other core areas in the region. In the Yakima core area some populations exhibit life history forms different from what they were historically. Migration between local populations and to and from spawning habitat is generally prevented or impeded by headwater storage dams on irrigation reservoirs, connectivity between tributaries and reservoirs, and within lower portions of spawning and rearing habitat and the mainstem Yakima River due to changed flow patterns, low instream flows, high water temperatures, and other habitat impediments. Currently, the connectivity in the Yakima Core area is truncated to the degree that not all populations are able to contribute gene flow to a functional metapopulation (USFWS 2015c, pp. C-6-7). Very little is known about the demographic status of the South Salmo Core Area. Irregular surveys conducted between 2005 and 2016, indicated multiple life stages and healthy presence.

Lower Snake Region

Demographic status is variable within the Lower Snake Region. Although trend data are lacking, several core areas in the Grande Ronde Basin and the Imnaha core area are thought to be stable. The upper Grande Ronde Core Area is the exception where population abundance is considered depressed. Wenaha, Little Minam, and Imnaha Rivers are strongholds (as mentioned above), as are most core areas in the Clearwater River basin. Most core areas contain populations that express both a resident and fluvial life history strategy. There is potential that some bull trout in the upper Wallowa River are adfluvial. There is potential for connectivity between core areas in the Grande Ronde basin, however conditions in FMO are limiting (USFWS 2015c, p. C-7).

Middle Snake Region

In the Middle Snake Region, core areas are distributed along both sides of the Snake River above Hells Canyon Dam. The Powder River and Pine Creek basins are in Oregon and Indian Creek and Wildhorse Creek are on the Idaho side of the Snake River. Demographic status of the core areas is poorest in the Powder River Core Area where populations are highly fragmented and severely depressed. The East Pine Creek population in the Pine-Indian-Wildhorse Creeks core area is likely the most abundant within the region. Populations in both core areas primarily express a resident life history strategy; however, some evidence suggests a migratory life history still exists in the Pine-Indian-Wildhorse Creeks core area. Connectivity is severely impaired in the Middle Snake Region. Dams, diversions and temperature barriers prevent movement among

populations and between core areas. Brownlee Dam isolates bull trout in Wildhorse Creek from other populations (USFWS 2015c, p. C-7).

Columbia Headwaters Recovery Unit

The Columbia Headwaters Recovery Unit (CHRU) includes western Montana, northern Idaho, and the northeastern corner of Washington. Major drainages include the Clark Fork River basin and its Flathead River contribution, the Kootenai River basin, and the Coeur d'Alene Lake basin. In this implementation plan for the CHRU we have slightly reorganized the structure from the 2002 Draft Recovery Plan, based on latest available science and fish passage improvements that have rejoined previously fragmented habitats. We now identify 35 bull trout core areas (compared to 47 in 2002) for this recovery unit. Fifteen of the 35 are referred to as "complex" core areas as they represent large interconnected habitats, each containing multiple spawning streams considered to host separate and largely genetically identifiable local populations. The 15 complex core areas contain the majority of individual bull trout and the bulk of the designated critical habitat (USFWS 2010).

However, somewhat unique to this recovery unit is the additional presence of 20 smaller core areas, each represented by a single local population. These "simple" core areas are found in remote glaciated headwater basins, often in Glacier National Park or federally-designated wilderness areas, but occasionally also in headwater valley bottoms. Many simple core areas are upstream of waterfalls or other natural barriers to fish migration. In these simple core areas bull trout have apparently persisted for thousands of years despite small populations and isolated existence. As such, simple core areas meet the criteria for core area designation and continue to be valued for their uniqueness, despite limitations of size and scope. Collectively, the 20 simple core areas contain less than 3 percent of the total bull trout core area habitat in the CHRU, but represent significant genetic and life history diversity (Meeuwig et al. 2010). Throughout this recovery unit implementation plan, we often separate our analyses to distinguish between complex and simple core areas, both in respect to threats as well as recovery actions (USFWS 2015d, pp. D-1-2).

In order to effectively manage the recovery unit implementation plan (RUIP) structure in this large and diverse landscape, the core areas have been separated into the following five natural geographic assemblages.

Upper Clark Fork Geographic Region

Starting at the Clark Fork River headwaters, the *Upper Clark Fork Geographic Region* comprises seven complex core areas, each of which occupies one or more major watersheds contributing to the Clark Fork basin (*i.e.*, Upper Clark Fork River, Rock Creek, Blackfoot River, Clearwater River and Lakes, Bitterroot River, West Fork Bitterroot River, and Middle Clark Fork River core areas) (USFWS 2015d, p. D-2).

Lower Clark Fork Geographic Region

The seven headwater core areas flow into the *Lower Clark Fork Geographic Region*, which comprises two complex core areas, Lake Pend Oreille and Priest Lake. Because of the systematic and jurisdictional complexity (three States and a Tribal entity) and the current degree of migratory fragmentation caused by five mainstem dams, the threats and recovery actions in

the Lake Pend Oreille (LPO) core area are very complex and are described in three parts. LPO-A is upstream of Cabinet Gorge Dam, almost entirely in Montana, and includes the mainstem Clark Fork River upstream to the confluence of the Flathead River as well as the portions of the lower Flathead River (e.g., Jocko River) on the Flathead Indian Reservation. LPO-B is the Pend Oreille lake basin proper and its tributaries, extending between Albeni Falls Dam downstream from the outlet of Lake Pend Oreille and Cabinet Gorge Dam just upstream of the lake; almost entirely in Idaho. LPO-C is the lower basin (i.e., lower Pend Oreille River), downstream of Albeni Falls Dam to Boundary Dam (1 mile upstream from the Canadian border) and bisected by Box Canyon Dam; including portions of Idaho, eastern Washington, and the Kalispel Reservation (USFWS 2015d, p. D-2).

Historically, and for current purposes of bull trout recovery, migratory connectivity among these separate fragments into a single entity remains a primary objective.

Flathead Geographic Region

The *Flathead Geographic Region* includes a major portion of northwestern Montana upstream of Kerr Dam on the outlet of Flathead Lake. The complex core area of Flathead Lake is the hub of this area, but other complex core areas isolated by dams are Hungry Horse Reservoir (formerly South Fork Flathead River) and Swan Lake. Within the glaciated basins of the Flathead River headwaters are 19 simple core areas, many of which lie in Glacier National Park or the Bob Marshall and Great Bear Wilderness areas and some of which are isolated by natural barriers or other features (USFWS 2015d, p. D-2).

Kootenai Geographic Region

To the northwest of the Flathead, in an entirely separate watershed, lies the *Kootenai Geographic Region*. The Kootenai is a uniquely patterned river system that originates in southeastern British Columbia, Canada. It dips, in a horseshoe configuration, into northwest Montana and north Idaho before turning north again to re-enter British Columbia and eventually join the Columbia River headwaters in British Columbia. The *Kootenai Geographic Region* contains two complex core areas (Lake Koocanusa and the Kootenai River) bisected since the 1970's by Libby Dam, and also a single naturally isolated simple core area (Bull Lake). Bull trout in both of the complex core areas retain strong migratory connections to populations in British Columbia (USFWS 2015d, p. D-3).

Coeur d'Alene Geographic Region

Finally, the *Coeur d'Alene Geographic Region* consists of a single, large complex core area centered on Coeur d'Alene Lake. It is grouped into the CHRU for purposes of physical and ecological similarity (adfluvial bull trout life history and nonanadromous linkage) rather than due to watershed connectivity with the rest of the CHRU, as it flows into the mid-Columbia River far downstream of the Clark Fork and Kootenai systems (USFWS 2015d, p. D-3).

Upper Snake Recovery Unit

The Upper Snake Recovery Unit includes portions of central Idaho, northern Nevada, and eastern Oregon. Major drainages include the Salmon River, Malheur River, Jarbidge River, Little Lost River, Boise River, Payette River, and the Weiser River. The Upper Snake Recovery

Unit contains 22 bull trout core areas within 7 geographic regions or major watersheds: Salmon River (10 core areas, 123 local populations), Boise River (2 core areas, 29 local populations), Payette River (5 core areas, 25 local populations), Little Lost River (1 core area, 10 local populations), Malheur River (2 core areas, 8 local populations), Jarbidge River (1 core area, 6 local populations), and Weiser River (1 core area, 5 local populations). The Upper Snake Recovery Unit includes a total of 206 local populations, with almost 60 percent being present in the Salmon River watershed (USFWS 2015e, p. E-1).

Three major bull trout life history expressions are present in the Upper Snake Recovery Unit, adfluvial³, fluvial⁴, and resident populations. Large areas of intact habitat exist primarily in the Salmon drainage, as this is the only drainage in the Upper Snake Recovery Unit that still flows directly into the Snake River; most other drainages no longer have direct connectivity due to irrigation uses or instream barriers. Bull trout in the Salmon basin share a genetic past with bull trout elsewhere in the Upper Snake Recovery Unit. Historically, the Upper Snake Recovery Unit is believed to have largely supported the fluvial life history form; however, many core areas are now isolated or have become fragmented watersheds, resulting in replacement of the fluvial life history with resident or adfluvial forms. The Weiser River, Squaw Creek, Pahsimeroi River, and North Fork Payette River core areas contain only resident populations of bull trout (USFWS 2015e, pp. E-1-2).

Salmon River

The Salmon River basin represents one of the few basins that are still free-flowing down to the Snake River. The core areas in the Salmon River basin do not have any major dams and a large extent (approximately 89 percent) is federally managed, with large portions of the Middle Fork Salmon River and Middle Fork Salmon River - Chamberlain core areas occurring within the Frank Church River of No Return Wilderness. Most core areas in the Salmon River basin contain large populations with many occupied stream segments. The Salmon River basin contains 10 of the 22 core areas in the Upper Snake Recovery Unit and contains the majority of the occupied habitat. Over 70 percent of occupied habitat in the Upper Snake Recovery Unit occurs in the Salmon River basin as well as 123 of the 206 local populations. Connectivity between core areas in the Salmon River basin is intact; therefore it is possible for fish in the mainstem Salmon to migrate to almost any Salmon River core area or even the Snake River. Connectivity within Salmon River basin core areas is mostly intact except for the Pahsimeroi River and portions of the Lemhi River. The Upper Salmon River, Lake Creek, and Opal Lake core areas contain adfluvial populations of bull trout, while most of the remaining core areas contain fluvial populations; only the Pahsimeroi contains strictly resident populations. Most core areas appear to have increasing or stable trends but trends are not known in the Pahsimeroi, Lake Creek, or Opal Lake core areas. The Idaho Department of Fish and Game reported trend data from 7 of the 10 core areas. This trend data indicated that populations were stable or increasing in the Upper Salmon River, Lemhi River, Middle Salmon River-Chamberlain, Little Lost River, and the South Fork Salmon River (IDFG 2005, 2008). Trends were stable or decreasing in the Little-Lower Salmon River, Middle Fork Salmon River, and the Middle Salmon River-Panther (IDFG 2005, 2008).

³ Adfluvial: Life history pattern of spawning and rearing in tributary streams and migrating to lakes or reservoirs to mature.

⁴ Fluvial: Life history pattern of spawning and rearing in tributary streams and migrating to larger rivers to mature.

Boise River

In the Boise River basin, two large dams are impassable barriers to upstream fish movement: Anderson Ranch Dam on the South Fork Boise River, and Arrowrock Dam on the mainstem Boise River. Fish in Anderson Ranch Reservoir have access to the South Fork Boise River upstream of the dam. Fish in Arrowrock Reservoir have access to the North Fork Boise River, Middle Fork Boise River, and lower South Fork Boise River. The Boise River basin contains 2 of the 22 core areas in the Upper Snake Recovery Unit. The core areas in the Boise River basin account for roughly 12 percent of occupied habitat in the Upper Snake Recovery Unit and contain 29 of the 206 local populations. Approximately 90 percent of both Arrowrock and Anderson Ranch core areas are federally owned; most lands are managed by the U.S. Forest Service, with some portions occurring in designated wilderness areas. Both the Arrowrock core area and the Anderson Ranch core area are isolated from other core areas. Both core areas contain fluvial bull trout that exhibit adfluvial characteristics and numerous resident populations. The Idaho Department of Fish and Game in 2014 determined that the Anderson Ranch core area had an increasing trend while trends in the Arrowrock core area is unknown (USFWS 2015e).

Payette River

The Payette River basin contains three major dams that are impassable barriers to fish: Deadwood Dam on the Deadwood River, Cascade Dam on the North Fork Payette River, and Black Canyon Reservoir on the Payette River. Only the Upper South Fork Payette River and the Middle Fork Payette River still have connectivity, the remaining core areas are isolated from each other due to dams. Both fluvial and adfluvial life history expression are still present in the Payette River basin but only resident populations are present in the Squaw Creek and North Fork Payette River core areas. The Payette River basin contains 5 of the 22 core areas and 25 of the 206 local populations in the recovery unit. Less than 9 percent of occupied habitat in the recovery unit is in this basin. Approximately 60 percent of the lands in the core areas are federally owned and the majority is managed by the U.S. Forest Service. Trend data are lacking and the current condition of the various core areas is unknown, but there is concern due to the current isolation of three (North Fork Payette River, Squaw Creek, Deadwood River) of the five core areas; the presence of only resident local populations in two (North Fork Payette River, Squaw Creek) of the five core areas; and the relatively low numbers present in the North Fork core area (USFWS 2015e, p. E-8).

Jarbidge River

The Jarbidge River core area contains two major fish barriers along the Bruneau River: the Buckaroo diversion and C. J. Strike Reservoir. Bull trout are not known to migrate down to the Snake River. There is one core area in the basin, with populations in the Jarbidge River; this watershed does not contain any barriers. Approximately 89 percent of the Jarbidge core area is federally owned. Most lands are managed by either the Forest Service or Bureau of Land Management. A large portion of the core area is within the Bruneau-Jarbidge Wilderness area. A tracking study has documented bull trout population connectivity among many of the local populations, in particular between West Fork Jarbidge River and Pine Creek. Movement between the East and West Fork Jarbidge River has also been documented; therefore both resident and fluvial populations are present. The core area contains six local populations and 3 percent of the occupied habitat in the recovery unit. Trend data are lacking within this core area (USFWS 2015e, p. E-9).

Little Lost River

The Little Lost River basin is unique in that the watershed is within a naturally occurring hydrologic sink and has no connectivity with other drainages. A small fluvial population of bull trout may still exist, but it appears that most populations are predominantly resident populations. There is one core area in the Little Lost basin, and approximately 89 percent of it is federally owned by either the U.S. Forest Service or Bureau of Land Management. The core area contains 10 local populations and less than 3 percent of the occupied habitat in the recovery unit. The current trend condition of this core area is likely stable, with most bull trout residing in Upper Sawmill Canyon (IDFG 2014).

Malheur River

The Malheur River basin contains major dams that are impassable to fish. The largest are Warm Springs Dam, impounding Warm Springs Reservoir on the mainstem Malheur River, and Agency Valley Dam, impounding Beulah Reservoir on the North Fork Malheur River. The dams result in two core areas that are isolated from each other and from other core areas. Local populations in the two core areas are limited to habitat in the upper watersheds. The Malheur River basin contains 2 of the 22 core areas and 8 of the 206 local populations in the recovery unit. Fluvial and resident populations are present in both core areas while adfluvial populations are present in the North Fork Malheur River. This basin contains less than 3 percent of the occupied habitat in the recovery unit, and approximately 60 percent of lands in the two core areas are federally owned. Trend data indicates that populations are declining in both core areas (USFWS 2015e, p. E-9).

Weiser River

The Weiser River basin contains local populations that are limited to habitat in the upper watersheds. The Weiser River basin contains only a single core area that consists of 5 of the 206 local populations in the recovery unit. Local populations occur in only three stream complexes in the upper watershed: 1) Upper Hornet Creek, 2) East Fork Weiser River, and 3) Upper Little Weiser River. These local populations include only resident life histories. This basin contains less than 2 percent of the occupied habitat in the recovery unit, and approximately 44 percent of lands are federally owned. Trend data from the Idaho Department of Fish and Game indicate that the populations in the Weiser core area are increasing (IDFG 2014) but it is considered vulnerable because local populations are isolated and likely do not express migratory life histories (USFWS 2015e, p.E-10).

St. Mary Recovery Unit

The Saint Mary Recovery Unit is located in northwest Montana east of the Continental Divide and includes the U.S. portions of the Saint Mary River basin, from its headwaters to the international boundary with Canada at the 49th parallel. The watershed and the bull trout population are linked to downstream aquatic resources in southern Alberta, Canada; the U.S. portion includes headwater spawning and rearing (SR) habitat in the tributaries and a portion of the FMO habitat in the mainstem of the Saint Mary River and Saint Mary lakes (Mogen and Kaeding 2001).

The Saint Mary Recovery Unit comprises four core areas; only one (Saint Mary River) is a complex core area with five described local bull trout populations (Divide, Boulder, Kennedy,

Otatso, and Lee Creeks). Roughly half of the linear extent of available FMO habitat in the mainstem Saint Mary system (between Saint Mary Falls at the upstream end and the downstream Canadian border) is comprised of Saint Mary and Lower Saint Mary Lakes, with the remainder in the Saint Mary River. The other three core areas (Slide Lakes, Cracker Lake, and Red Eagle Lake) are simple core areas. Slide Lakes and Cracker Lake occur upstream of seasonal or permanent barriers and are comprised of genetically isolated single local bull trout populations, wholly within Glacier National Park, Montana. In the case of Red Eagle Lake, physical isolation does not occur, but consistent with other lakes in the adjacent Columbia Headwaters Recovery Unit, there is likely some degree of spatial separation from downstream Saint Mary Lake. As noted, the extent of isolation has been identified as a research need (USFWS 2015f, p. F-1). Bull trout in the Saint Mary River complex core area are documented to exhibit primarily the migratory fluvial life history form (Mogen and Kaeding 2005a, 2005b), but there is doubtless some occupancy (though less well documented) of Saint Mary Lakes, suggesting a partly adfluvial adaptation. Since lake trout and northern pike are both native to the Saint Mary River system (headwaters of the South Saskatchewan River drainage draining to Hudson Bay), the conventional wisdom is that these large piscivores historically outcompeted bull trout in the lacustrine environment (Donald and Alger 1993, Martinez et al. 2009), resulting in a primarily fluvial niche and existence for bull trout in this system. This is an untested hypothesis and additional research into this aspect is needed (USFWS 2015f, p. F-3).

Bull trout populations in the simple core areas of the three headwater lake systems (Slide, Cracker, and Red Eagle Lakes) are, by definition, adfluvial; there are also resident life history components in portions of the Saint Mary River system such as Lower Otatso Creek (Mogen and Kaeding 2005a), further exemplifying the overall life history diversity typical of bull trout. Mogen and Kaeding (2001) reported that bull trout continue to inhabit nearly all suitable habitats accessible to them in the Saint Mary River basin in the United States. The possible exception is portions of Divide Creek, which appears to be intermittently occupied despite a lack of permanent migratory barriers, possibly due to low population size and erratic year class production (USFWS 2015f, p. F-3).

It should be noted that bull trout are found in minor portions of two additional U.S. watersheds (Belly and Waterton rivers) that were once included in the original draft recovery plan (USFWS 2002) but are no longer considered core areas in the final recovery plan (USFWS 2015) and are not addressed in that document. In Alberta, Canada, the Saint Mary River bull trout population is considered at “high risk,” while the Belly River is rated as “at risk” (ACA 2009). In the Belly River drainage, which enters the South Saskatchewan system downstream of the Saint Mary River in Alberta, some bull trout spawning is known to occur on either side of the international boundary. These waters are in the drainage immediately west of the Saint Mary River headwaters. However, the U.S. range of this population constitutes only a minor headwater migratory SR segment of an otherwise wholly Canadian population, extending less than 1 mile (0.6 km) into backcountry waters of Glacier National Park. The Belly River population is otherwise totally dependent on management within Canadian jurisdiction, with no natural migratory connection to the Saint Mary (USFWS 2015f, p. F-3).

Current status of bull trout in the Saint Mary River core area (U.S.) is considered strong (Mogen 2013). Migratory bull trout redd counts are conducted annually in the two major SR streams,

Boulder and Kennedy creeks. Boulder Creek redd counts have ranged from 33 to 66 in the past decade, with the last 4 counts all 53 or higher. Kennedy Creek redd counts are less robust, ranging from 5 to 25 over the last decade, with a 2014 count of 20 (USFWS 2015f, p. F-3). Generally, the demographic status of the Saint Mary River core area is believed to be good, with the exception of the Divide Creek local population. In this local population, there is evidence that a combination of ongoing habitat manipulation (Smillie and Ellerbroek 1991, F-5 NPS 1992) resulting in occasional historical passage issues, combined with low and erratic recruitment (DeHaan et al. 2011) has caused concern for the continuing existence of the local population. While less is known about the demographic status of the three simple cores where redd counts are not conducted, all three appear to be self-sustaining and fluctuating within known historical population demographic bounds. Of the three simple core areas, demographic status in Slide Lakes and Cracker Lake appear to be functioning appropriately, but the demographic status in Red Eagle Lake is less well documented and believed to be less robust (USFWS 2015f, p. F-3).

Reasons for Listing

Bull trout distribution, abundance, and habitat quality have declined rangewide (Bond 1992, pp. 2-3; Schill 1992, p. 42; Thomas 1992, entire; Ziller 1992, entire; Rieman and McIntyre 1993, p. 1; Newton and Pribyl 1994, pp. 4-5; McPhail and Baxter 1996, p. 1). Several local extirpations have been documented, beginning in the 1950s (Rode 1990, pp. 26-32; Ratliff and Howell 1992, entire; Donald and Alger 1993, entire; Goetz 1994, p. 1; Newton and Pribyl 1994, pp. 8-9; Light et al. 1996, pp. 6-7; Buchanan et al. 1997, p. 15; WDFW 1998, pp. 2-3). Bull trout were extirpated from the southernmost portion of their historic range, the McCloud River in California, around 1975 (Rode 1990, p. 32). Bull trout have been functionally extirpated (i.e., few individuals may occur there but do not constitute a viable population) in the Coeur d'Alene River basin in Idaho and in the Lake Chelan and Okanogan River basins in Washington (USFWS 1998, pp. 31651-31652).

These declines result from the combined effects of habitat degradation and fragmentation, the blockage of migratory corridors; poor water quality, angler harvest and poaching, entrainment (process by which aquatic organisms are pulled through a diversion or other device) into diversion channels and dams, and introduced nonnative species. Specific land and water management activities that depress bull trout populations and degrade habitat include the effects of dams and other diversion structures, forest management practices, livestock grazing, agriculture, agricultural diversions, road construction and maintenance, mining, and urban and rural development (Beschta et al. 1987, entire; Chamberlain et al. 1991, entire; Furniss et al. 1991, entire; Meehan 1991, entire; Nehlsen et al. 1991, entire; Sedell and Everest 1991, entire; Craig and Wissmar 1993pp, 18-19; Henjum et al. 1994, pp. 5-6; McIntosh et al. 1994, entire; Wissmar et al. 1994, entire; MBTSG 1995a, p. 1; MBTSG 1995b, pp. i-ii; MBTSG 1995c, pp. i-ii; MBTSG 1995d, p. 22; MBTSG 1995e, p. i; MBTSG 1996a, p. i-ii; MBTSG 1996b, p. i; MBTSG 1996c, p. i; MBTSG 1996d, p. i; MBTSG 1996e, p. i; MBTSG 1996f, p. 11; Light et al. 1996, pp. 6-7; USDA and USDI 1995, p. 2).

Emerging Threats

Climate Change

Climate change was not addressed as a known threat when bull trout was listed. The 2015 bull trout recovery plan and RUIPs summarize the threat of climate change and acknowledges that some extant bull trout core area habitats will likely change (and may be lost) over time due to anthropogenic climate change effects, and use of best available information will ensure future conservation efforts that offer the greatest long-term benefit to sustain bull trout and their required coldwater habitats (USFWS 2015, p. vii, and pp. 17-20, USFWS 2015a-f).

Global climate change and the related warming of global climate have been well documented (IPCC 2007, entire; ISAB 2007, entire; Combes 2003, entire). Evidence of global climate change/warming includes widespread increases in average air and ocean temperatures and accelerated melting of glaciers, and rising sea level. Given the increasing certainty that climate change is occurring and is accelerating (IPCC 2007, p. 253; Battin et al. 2007, p. 6720), we can no longer assume that climate conditions in the future will resemble those in the past.

Patterns consistent with changes in climate have already been observed in the range of many species and in a wide range of environmental trends (ISAB 2007, entire; Hari et al. 2006, entire; Rieman et al. 2007, entire). In the northern hemisphere, the duration of ice cover over lakes and rivers has decreased by almost 20 days since the mid-1800's (Magnuson et al. 2000, p. 1743). The range of many species has shifted poleward and elevationally upward. For cold-water associated salmonids in mountainous regions, where their upper distribution is often limited by impassable barriers, an upward thermal shift in suitable habitat can result in a reduction in range, which in turn can lead to a population decline (Hari et al. 2006, entire).

In the Pacific Northwest, most models project warmer air temperatures and increases in winter precipitation and decreases in summer precipitation. Warmer temperatures will lead to more precipitation falling as rain rather than snow. As the seasonal amount of snow pack diminishes, the timing and volume of stream flow are likely to change and peak river flows are likely to increase in affected areas. Higher air temperatures are also likely to increase water temperatures (ISAB 2007, pp. 15-17). For example, stream gauge data from western Washington over the past 5 to 25 years indicate a marked increasing trend in water temperatures in most major rivers. Climate change has the potential to profoundly alter the aquatic ecosystems upon which the bull trout depends via alterations in water yield, peak flows, and stream temperature, and an increase in the frequency and magnitude of catastrophic wildfires in adjacent terrestrial habitats (Bisson et al. 2003, pp 216-217).

All life stages of the bull trout rely on cold water. Increasing air temperatures are likely to impact the availability of suitable cold water habitat. For example, ground water temperature is generally correlated with mean annual air temperature, and has been shown to strongly influence the distribution of other chars. Ground water temperature is linked to bull trout selection of spawning sites, and has been shown to influence the survival of embryos and early juvenile rearing of bull trout (Baxter 1997, p. 82). Increases in air temperature are likely to be reflected in increases in both surface and groundwater temperatures.

Climate change is likely to affect the frequency and magnitude of fires, especially in warmer drier areas such as are found on the eastside of the Cascade Mountains. Bisson et al. (2003, pp. 216-217) note that the forest that naturally occurred in a particular area may or may not be the forest that will be responding to the fire regimes of an altered climate. In several studies related to the effect of large fires on bull trout populations, bull trout appear to have adapted to past fire disturbances through mechanisms such as dispersal and plasticity. However, as stated earlier, the future may well be different than the past and extreme fire events may have a dramatic effect on bull trout and other aquatic species, especially in the context of continued habitat loss, simplification and fragmentation of aquatic systems, and the introduction and expansion of exotic species (Bisson et al. 2003, pp. 218-219).

Migratory bull trout can be found in lakes, large rivers and marine waters. Effects of climate change on lakes are likely to impact migratory adfluvial bull trout that seasonally rely upon lakes for their greater availability of prey and access to tributaries. Climate-warming impacts to lakes will likely lead to longer periods of thermal stratification and coldwater fish such as adfluvial bull trout will be restricted to these bottom layers for greater periods of time. Deeper thermoclines resulting from climate change may further reduce the area of suitable temperatures in the bottom layers and intensify competition for food (Shuter and Meisner 1992, p. 11). Bull trout require very cold water for spawning and incubation. Suitable spawning habitat is often found in accessible higher elevation tributaries and headwaters of rivers. However, impacts on hydrology associated with climate change are related to shifts in timing, magnitude and distribution of peak flows that are also likely to be most pronounced in these high elevation stream basins (Battin et al. 2007, p. 6720). The increased magnitude of winter peak flows in high elevation areas is likely to impact the location, timing, and success of spawning and incubation for the bull trout and Pacific salmon species. Although lower elevation river reaches are not expected to experience as severe an impact from alterations in stream hydrology, they are unlikely to provide suitably cold temperatures for bull trout spawning, incubation and juvenile rearing.

As climate change progresses and stream temperatures warm, thermal refugia will be critical to the persistence of many bull trout populations. Thermal refugia are important for providing bull trout with patches of suitable habitat during migration through or to make feeding forays into areas with greater than optimal temperatures.

There is still a great deal of uncertainty associated with predictions relative to the timing, location, and magnitude of future climate change. It is also likely that the intensity of effects will vary by region (ISAB 2007, p 7) although the scale of that variation may exceed that of States. For example, several studies indicate that climate change has the potential to impact ecosystems in nearly all streams throughout the State of Washington (ISAB 2007, p. 13; Battin et al. 2007, p. 6722; Rieman et al. 2007, pp. 1558-1561). In streams and rivers with temperatures approaching or at the upper limit of allowable water temperatures, there is little if any likelihood that bull trout will be able to adapt to or avoid the effects of climate change/warming. There is little doubt that climate change is and will be an important factor affecting bull trout distribution. As its distribution contracts, patch size decreases and connectivity is truncated, bull trout populations that may be currently connected may face increasing isolation, which could accelerate the rate of local extinction beyond that resulting from changes in stream temperature

alone (Rieman et al. 2007, pp. 1559-1560). Due to variations in land form and geographic location across the range of the bull trout, it appears that some populations face higher risks than others. Bull trout in areas with currently degraded water temperatures and/or at the southern edge of its range may already be at risk of adverse impacts from current as well as future climate change.

The ability to assign the effects of gradual global climate change to bull trout or to a specific location on the ground is beyond our technical capabilities at this time.

Conservation

Conservation Needs

The 2015 recovery plan for bull trout established the primary strategy for recovery of bull trout in the coterminous United States: 1) conserve bull trout so that they are geographically widespread across representative habitats and demographically stable in six recovery units; 2) effectively manage and ameliorate the primary threats in each of six recovery units at the core area scale such that bull trout are not likely to become endangered in the foreseeable future; 3) build upon the numerous and ongoing conservation actions implemented on behalf of bull trout since their listing in 1999, and improve our understanding of how various threat factors potentially affect the species; 4) use that information to work cooperatively with our partners to design, fund, prioritize, and implement effective conservation actions in those areas that offer the greatest long-term benefit to sustain bull trout and where recovery can be achieved; and 5) apply adaptive management principles to implementing the bull trout recovery program to account for new information (USFWS 2015, p. v.).

Information presented in prior draft recovery plans published in 2002 and 2004 (USFWS 2002a, 2004) have served to identify recovery actions across the range of the species and to provide a framework for implementing numerous recovery actions by our partner agencies, local working groups, and others with an interest in bull trout conservation.

The 2015 recovery plan (USFWS 2015) integrates new information collected since the 1999 listing regarding bull trout life history, distribution, demographics, conservation successes, etc., and integrates and updates previous bull trout recovery planning efforts across the range of the single DPS listed under the Endangered Species Act of 1973, as amended (16 U.S.C. 1531 *et seq.*) (Act).

The Service has developed a recovery approach that: 1) focuses on the identification of and effective management of known and remaining threat factors to bull trout in each core area; 2) acknowledges that some extant bull trout core area habitats will likely change (and may be lost) over time; and 3) identifies and focuses recovery actions in those areas where success is likely to meet our goal of ensuring the certainty of conservation of genetic diversity, life history features, and broad geographical representation of remaining bull trout populations so that the protections of the Act are no longer necessary (USFWS 2015, p. 45-46).

To implement the recovery strategy, the 2015 recovery plan establishes categories of recovery actions for each of the six Recovery Units (USFWS 2015, p. 50-51):

1. Protect, restore, and maintain suitable habitat conditions for bull trout.
2. Minimize demographic threats to bull trout by restoring connectivity or populations where appropriate to promote diverse life history strategies and conserve genetic diversity.

3. Prevent and reduce negative effects of nonnative fishes and other nonnative taxa on bull trout.
4. Work with partners to conduct research and monitoring to implement and evaluate bull trout recovery activities, consistent with an adaptive management approach using feedback from implemented, site-specific recovery tasks, and considering the effects of climate change.

Bull trout recovery is based on a geographical hierarchical approach. Bull trout are listed as a single DPS within the five-state area of the coterminous United States. The single DPS is subdivided into six biologically-based recovery units: 1) Coastal Recovery Unit; 2) Klamath Recovery Unit; 3) Mid-Columbia Recovery Unit; 4) Upper Snake Recovery Unit; 5) Columbia Headwaters Recovery Unit; and 6) Saint Mary Recovery Unit (USFWS 2015, p. 23). A viable recovery unit should demonstrate that the three primary principles of biodiversity have been met: representation (conserving the genetic makeup of the species); resiliency (ensuring that each population is sufficiently large to withstand stochastic events); and redundancy (ensuring a sufficient number of populations to withstand catastrophic events) (USFWS 2015, p. 33). Each of the six recovery units contain multiple bull trout core areas, 116 total, which are non-overlapping watershed-based polygons, and each core area includes one or more local populations. Currently there are 109 occupied core areas, which comprise 611 local populations (USFWS 2015, p. 3). There are also six core areas where bull trout historically occurred but are now extirpated, and one research needs area where bull trout were known to occur historically, but their current presence and use of the area are uncertain (USFWS 2015, p. 3). Core areas can be further described as complex or simple (USFWS 2015, p. 3-4). Complex core areas contain multiple local bull trout populations, are found in large watersheds, have multiple life history forms, and have migratory connectivity between spawning and rearing habitat and FMO habitats. Simple core areas are those that contain one bull trout local population. Simple core areas are small in scope, isolated from other core areas by natural barriers, and may contain unique genetic or life history adaptations.

A local population is a group of bull trout that spawn within a particular stream or portion of a stream system (USFWS 2015, p. 73). A local population is considered to be the smallest group of fish that is known to represent an interacting reproductive unit. For most waters where specific information is lacking, a local population may be represented by a single headwater tributary or complex of headwater tributaries. Gene flow may occur between local populations (e.g., those within a core population), but is assumed to be infrequent compared with that among individuals within a local population.

Recovery Units and Local Populations

The final recovery plan (USFWS 2015) designates six bull trout recovery units as described above. These units replace the 5 interim recovery units previously identified (USFWS 1999). The Service will address the conservation of these final recovery units in our section 7(a)(2) analysis for proposed Federal actions. The recovery plan (USFWS 2015), identified threats and factors affecting the bull trout within these units. A detailed description of recovery implementation for each recovery unit is provided in separate RUIPs (USFWS 2015a-f), which identify conservation actions and recommendations needed for each core area, forage/ migration/

overwinter areas, historical core areas, and research needs areas. Each of the following recovery units (below) is necessary to maintain the bull trout's distribution, as well as its genetic and phenotypic diversity, all of which are important to ensure the species' resilience to changing environmental conditions.

Coastal Recovery Unit

The coastal recovery unit implementation plan describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the unit (USFWS 2015a). The Coastal Recovery Unit is located within western Oregon and Washington. The Coastal Recovery Unit is divided into three regions: Puget Sound, Olympic Peninsula, and the Lower Columbia River Regions. This recovery unit contains 20 core areas comprising 84 local populations and a single potential local population in the historic Clackamas River core area where bull trout had been extirpated and were reintroduced in 2011, and identified four historically occupied core areas that could be re-established (USFWS 2015, pg. 47; USFWS 2015a, p. A-2). Core areas within Puget Sound and the Olympic Peninsula currently support the only anadromous local populations of bull trout. This recovery unit also contains ten shared FMO habitats which are outside core areas and allows for the continued natural population dynamics in which the core areas have evolved (USFWS 2015a, p. A-5). There are four core areas within the Coastal Recovery Unit that have been identified as current population strongholds: Lower Skagit, Upper Skagit, Quinault River, and Lower Deschutes River (USFWS 2015, p.79). These are the most stable and abundant bull trout populations in the recovery unit. The current condition of the bull trout in this recovery unit is attributed to the adverse effects of climate change, loss of functioning estuarine and nearshore marine habitats, development and related impacts (e.g., flood control, floodplain disconnection, bank armoring, channel straightening, loss of instream habitat complexity), agriculture (e.g., diking, water control structures, draining of wetlands, channelization, and the removal of riparian vegetation, livestock grazing), fish passage (e.g., dams, culverts, instream flows) residential development, urbanization, forest management practices (e.g., timber harvest and associated road building activities), connectivity impairment, mining, and the introduction of non-native species. Conservation measures or recovery actions implemented include relicensing of major hydropower facilities that have provided upstream and downstream fish passage or complete removal of dams, land acquisition to conserve bull trout habitat, floodplain restoration, culvert removal, riparian revegetation, levee setbacks, road removal, and projects to protect and restore important nearshore marine habitats.

Klamath Recovery Unit

The Klamath recovery unit implementation plan describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the unit (USFWS 2015b). The Klamath Recovery Unit is located in southern Oregon and northwestern California. The Klamath Recovery Unit is the most significantly imperiled recovery unit, having experienced considerable extirpation and geographic contraction of local populations and declining demographic condition, and natural re-colonization is constrained by dispersal barriers and presence of nonnative brook trout (USFWS 2015, p. 39). This recovery unit currently contains three core areas and eight local populations (USFWS 2015, p. 47; USFWS 2015b, p. B-1). Nine historic local populations of bull trout have become extirpated (USFWS 2015b, p.

B-1). All three core areas have been isolated from other bull trout populations for the past 10,000 years (USFWS 2015b, p. B-3). The current condition of the bull trout in this recovery unit is attributed to the adverse effects of climate change, habitat degradation and fragmentation, past and present land use practices, agricultural water diversions, nonnative species, and past fisheries management practices. Conservation measures or recovery actions implemented include removal of nonnative fish (e.g., brook trout, brown trout, and hybrids), acquiring water rights for instream flows, replacing diversion structures, installing fish screens, constructing bypass channels, installing riparian fencing, culver replacement, and habitat restoration.

Mid-Columbia Recovery Unit

The Mid-Columbia recovery unit implementation plan describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the unit (USFWS 2015c). The Mid-Columbia Recovery Unit is located within eastern Washington, eastern Oregon, and portions of central Idaho. The Mid-Columbia Recovery Unit is divided into four geographic regions: Lower Mid-Columbia, Upper Mid-Columbia, Lower Snake, and Mid-Snake Geographic Regions. This recovery unit contains 24 occupied core areas comprising 142 local populations, two historically occupied core areas, one research needs area, and seven FMO habitats (USFWS 2015, pg. 47; USFWS 2015c, p. C-1–4). The current condition of the bull trout in this recovery unit is attributed to the adverse effects of climate change, agricultural practices (e.g. irrigation, water withdrawals, livestock grazing), fish passage (e.g. dams, culverts), nonnative species, forest management practices, and mining. Conservation measures or recovery actions implemented include road removal, channel restoration, mine reclamation, improved grazing management, removal of fish barriers, and instream flow requirements.

Columbia Headwaters Recovery Unit

The Columbia headwaters recovery unit implementation plan describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the unit (USFWS 2015d, entire). The Columbia Headwaters Recovery Unit is located in western Montana, northern Idaho, and the northeastern corner of Washington. The Columbia Headwaters Recovery Unit is divided into five geographic regions: Upper Clark Fork, Lower Clark Fork, Flathead, Kootenai, and Coeur d'Alene Geographic Regions (USFWS 2015d, pp. D-2 – D-4). This recovery unit contains 35 bull trout core areas; 15 of which are complex core areas as they represent larger interconnected habitats and 20 simple core areas as they are isolated headwater lakes with single local populations. The 20 simple core areas are each represented by a single local population, many of which may have persisted for thousands of years despite small populations and isolated existence (USFWS 2015d, p. D-1). Fish passage improvements within the recovery unit have reconnected some previously fragmented habitats (USFWS 2015d, p. D-1), while others remain fragmented. Unlike the other recovery units in Washington, Idaho and Oregon, the Columbia Headwaters Recovery Unit does not have any anadromous fish overlap. Therefore, bull trout within the Columbia Headwaters Recovery Unit do not benefit from the recovery actions for salmon (USFWS 2015d, p. D-41). The current condition of the bull trout in this recovery unit is attributed to the adverse effects of climate change, mostly historical mining and contamination by heavy metals, expanding populations of nonnative fish predators and competitors, modified instream flows, migratory barriers (e.g., dams), habitat fragmentation, forest practices (e.g., logging, roads), agriculture practices (e.g. irrigation, livestock grazing), and residential development. Conservation measures or recovery

actions implemented include habitat improvement, fish passage, and removal of nonnative species.

Upper Snake Recovery Unit

The Upper Snake recovery unit implementation plan describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the unit (USFWS 2015e, entire). The Upper Snake Recovery Unit is located in central Idaho, northern Nevada, and eastern Oregon. The Upper Snake Recovery Unit is divided into seven geographic regions: Salmon River, Boise River, Payette River, Little Lost River, Malheur River, Jarbidge River, and Weiser River. This recovery unit contains 22 core areas and 207 local populations (USFWS 2015, p. 47), with almost 60 percent being present in the Salmon River Region. The current condition of the bull trout in this recovery unit is attributed to the adverse effects of climate change, dams, mining, forest management practices, nonnative species, and agriculture (e.g., water diversions, grazing). Conservation measures or recovery actions implemented include instream habitat restoration, instream flow requirements, screening of irrigation diversions, and riparian restoration.

St. Mary Recovery Unit

The St. Mary recovery unit implementation plan describes the threats to bull trout and the site-specific management actions necessary for recovery of the species within the unit (USFWS 2015f). The Saint Mary Recovery Unit is located in Montana but is heavily linked to downstream resources in southern Alberta, Canada. Most of the Saskatchewan River watershed which the St. Mary flows into is located in Canada. The United States portion includes headwater spawning and rearing habitat and the upper reaches of FMO habitat. This recovery unit contains four core areas, and seven local populations (USFWS 2015f, p. F-1) in the U.S. Headwaters. The current condition of the bull trout in this recovery unit is attributed primarily to the outdated design and operations of the Saint Mary Diversion operated by the Bureau of Reclamation (e.g., entrainment, fish passage, instream flows), and, to a lesser extent habitat impacts from development and nonnative species.

Tribal Conservation Activities

Many Tribes throughout the range of the bull trout are participating on bull trout conservation working groups or recovery teams in their geographic areas of interest. Some tribes are also implementing projects which focus on bull trout or that address anadromous fish but benefit bull trout (e.g., habitat surveys, passage at dams and diversions, habitat improvement, and movement studies).

LITERATURE CITED

- [ACA] Alberta Sustainable Resource Development and Alberta Conservation Association. 2009. Status of the bull trout (*Salvelinus confluentus*) in Alberta: Update 2009. Alberta Sustainable Resource Development. Wildlife Status Report No. 39 (Update 2009). Edmonton, Alberta.

- Ardren, W. R., P. W. DeHaan, C. T. Smith, E. B. Taylor, R. Leary, C. C. Kozfkay, L. Godfrey, M. Diggs, W. Fredenberg, and J. Chan. 2011. Genetic structure, evolutionary history, and conservation units of bull trout in the coterminous United States. *Transactions of the American Fisheries Society* 140:506-525. 22 pp.
- Battin, J., M.W. Wiley, M.H. Ruckelshaus, R.N. Palmer, E. Korb, K.K. Bartz, and H. Imaki. 2007. Projected impacts of climate change on salmon habitat restoration. *Proceedings of the National Academy of Sciences of the United States of America* 104(16):6720-6725. 6 pp.
- Baxter, C.V. 2002. Fish movement and assemblage dynamics in a Pacific Northwest riverscape. Doctoral dissertation. Oregon State University, Corvallis, OR. 174 pp.
- Baxter, J. S. 1997. Aspects of the reproductive ecology of bull trout in the Chowade River, British Columbia. Master's thesis. University of British Columbia, Vancouver. 110 pp.
- Beauchamp, D.A., and J.J. VanTassell. 2001. Modeling seasonal trophic interactions of adfluvial bull trout in Lake Billy Chinook, Oregon. *Transactions of the American Fisheries Society* 130:204-216. 13 pp.
- Behnke, R.J. 2002. Trout and Salmon of North America; Chapter: Bull Trout. Free Press, Simon and Shuster, Inc. N.Y., N.Y. Pp. 293-299.
- Beschta, R.L., R.E. Bilby, G.W. Brown, L.B. Holtby, and T.D. Hofstra. 1987. Stream temperature and aquatic habitat: fisheries and forestry interactions. Pages 191-232 in E.D. Salo and T.W. Cundy (eds). *Streamside Management Forestry and Fisheries Interactions*. Institute of Forest Resources, University of Washington, Seattle, Washington, Contribution No. 57. 46 pp.
- Bisson, P.A., B.E. Rieman, C. Luce, P.F. Hessburg, D.C. Lee, J.L. Kershner, G.H. Reeves, and R.E. Gresswell. 2003. Fire and aquatic ecosystems of the western USA: Current knowledge and key questions. *Forest Ecology and Management*. 178 (2003) 213-229. 17 pp.
- Boag, T.D. 1987. Food habits of bull char, *Salvelinus confluentus*, and rainbow trout, *Salmo gairdneri*, coexisting in a foothills stream in northern Alberta. *Canadian Field-Naturalist* 101(1): 56-62. 6 pp.
- Bond, C.E. 1992. Notes on the nomenclature and distribution of the bull trout and the effects of human activity on the species. Pages 1-4 in Howell, P.J. and D.V. 4 pp.
- Bonneau, J.L. and D.L. Scarnecchia. 1996. Distribution of juvenile bull trout in a thermal gradient of a plunge pool in Granite Creek, Idaho. *Transactions of the American Fisheries Society* 125: 628-630. 3 pp.
- Brenkman, S.J. and S.C. Corbett. 2005. Extent of Anadromy in Bull Trout and Implications for Conservation of a Threatened Species. *North American Journal of Fisheries Management*. 25:1073-1081. 9 pp.

- Brewin, P.A. and M. K. Brewin. 1997. Distribution Maps for Bull Trout in Alberta. Pages 206-216 in Mackay, W.C., M.K. Brewin and M. Monita. Friends of the bull Trout Conference Proceedings. 10 pp.
- Buchanan, D.V., and S.V. Gregory. 1997. Development of water temperature standards to protect and restore habitat for bull trout and other cold water species in Oregon. Mackay, W.C., Pp. 119-126
- Buchanan, D.V., M.L. Hanson, and R.M. Hooton. 1997. Status of Oregon's bull trout. Oregon Department of Fish and Wildlife. 168 pp.
- Buktenica, M. W., D. K. Hering, S. F. Girdner, B. D. Mahoney, and B. D. Rosenlund. 2013. Eradication of nonnative brook trout with electrofishing and antimycin-A and the response of a remnant bull trout population. *North American Journal of Fisheries Management* 33:117-129.
- Burkey, T.V. 1989. Extinction in nature reserves: the effect of fragmentation and the importance of migration between reserve fragments. *Oikos* 55:75-81. 7 pp.
- Burkey, T.V. 1995. Extinction rates in archipelagoes: Implications for populations in fragmented habitats. *Conservation Biology* 9: 527-541. 16 pp.
- Cavender, T. M. 1978. Taxonomy and distribution of the bull trout, *Salvelinus confluentus* (Suckley), from the American Northwest. *California Fish and Game* 64: 139-174. 19 pp.
- Chamberlain, T. W., R. D. Harr, and F. H. Everest. 1991. Timber harvesting, silviculture and watershed processes. Pages 181-205 in W. R. Meehan (ed). Influences of forest and rangeland management on salmonid fishes and their habitats. *American Fisheries Society Special Publication* 19. 26 pp.
- Combes, S. 2003. Protecting freshwater ecosystems in the face of global climate change. In: Hansen LJ et al. (eds) *Buying time: a user's manual for building resistance and resilience to climate change in natural systems*. WWF, Washington, UDA. Pp. 175-214. 44 pp.
- Costello, A.B., T.E. Down, S.M. Pollard, C.J. Pacas, and E.B. Taylor. 2003. The influence of history and contemporary stream hydrology on the evolution of genetic diversity within species: an examination of microsatellite DNA variation in bull trout, *Salvelinus confluentus* (Pisces: Salmonidae). *Evolution*. 57(2):328-344. 17 pp.
- Craig, S.D., and R.C. Wissmar. 1993. Habitat conditions influencing a remnant bull trout spawning population, Gold Creek, Washington (draft report). Fisheries Research Institute, University of Washington. Seattle, Washington. 47 pp.
- Dambacher, J. M., M. W. Buktenica, and G. L. Larson. 1992. Distribution, abundance, and habitat utilization of bull trout and brook trout in Sun Creek, Crater Lake National Park, Oregon. Proceedings of the Gearhart Mountain Bull Trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon.

- DeHaan, P., M. Diggs, and J. VonBargen. 2011. Genetic analysis of bull trout in the Saint Mary River System. U.S. Fish and Wildlife Service. Abernathy Fish Technology Center, Longview, Washington.
- Donald, D.B. and D.J. Alger. 1993. Geographic distribution, species displacement, and niche overlap for lake trout and bull trout in mountain lakes. *Canadian Journal of Zoology* 71: 238-247. 10 pp.
- Dunham, J.B. and B.E. Rieman. 1999. Metapopulation structure of bull trout: Influences of physical, biotic, and geometrical landscape characteristics. *Ecological Applications* 9:642-655. 15 pp.
- Dunham, J., B. Rieman, and G. Chandler. 2003. Influences of temperature and environmental variables on the distribution of bull trout within streams at the southern margin of its range. *North American Journal of Fisheries Management* 23:894-905. 11 pp.
- Dunham, J., C. Baxter, K. Fausch, W. Fredenberg, S. Kitano, I. Koizumi, K. Morita, T. Nakamura, B. Rieman, K. Savvaitova, J. Stanford, E. Taylor, and S. Yamamoto. 2008. Evolution, ecology, and conservation of Dolly Varden, white-spotted char, and bull trout. *Fisheries* 33:537-550.
- Fishbase 2015. <http://www.fishbase.org/Summary/SpeciesSummary.php?ID=2690&AT=bull+trout> 2pp.
- Fraley, J.J., and B.B. Shepard. 1989. Life history, ecology and population status of migratory bull trout (*Salvelinus confluentus*) in the Flathead Lake and River System, Montana. *Northwest Science* 63(4):133-143.
- Frissell, C.A. 1999. An ecosystem approach to habitat conservation for bull trout: groundwater and surface water protection. Open File Report Number 156-99. Flathead Lake Biological Station, University of Montana, Polson, MT, 46 pp.
- Furniss, M.J., T.D. Roelofs, and C.S. Yee. 1991. Road construction and maintenance. *American Fisheries Society Special Publication* 19:297-323. 14 pp.
- Gilbert C. H. 1897. The fishes of the Klamath Basin. *Bulletin of the U.S. Fish Commission* 17:1-13.
- Goetz, F. 1989. Biology of the bull trout, *Salvelinus confluentus*, a literature review. Willamette National Forest. Eugene, Oregon. 60 pp.
- Goetz, F. 1994. Distribution and juvenile ecology of bull trout (*Salvelinus confluentus*) in the Cascade Mountains. M.S. thesis. Oregon State University, Corvallis. 190 pp.
- Goetz, F., E. Jeanes, and E. Beamer. 2004. Bull trout in the nearshore. Preliminary draft. U.S. Army Corps of Engineers, Seattle, Washington, June, 2004, 396 pp.

- Haas, G. R., and J. D. McPhail. 1991. Systematics and distributions of Dolly Varden (*Salvelinus malma*) and bull trout (*Salvelinus confluentus*) in North America. *Can. J. Fish. Aquat. Sci.* 48: 2191-2211. 21 pp.
- Hartill, T. and S. Jacobs. 2007. Distribution and abundance of bull trout in the Sprague River (Upper Klamath Basin), 2006. Oregon Department of Fish and Wildlife. Corvallis, Oregon.
- Hari, R. E., D. M. Livingstone, R. Siber, P. Burkhardt-Holm, and H. Guttinger. 2006. Consequences of climatic change for water temperature and brown trout populations in alpine rivers and streams. *Global Change Biology* 12:10-26. 17 pp.
- Henjum, M.G., J.R. Karr, D.L. Bottom, D.A. Perry, J.C. Bednarz, S.G. Wright, S.A. Beckwitt, and E. Beckwitt. 1994. Interim protection for late-successional forests, fisheries, and watersheds. National forests east of the Cascade Crest, Oregon, and Washington. A report to the Congress and President of the United States Eastside Forests Scientific Society Panel. American Fisheries Society, American Ornithologists Union Incorporated, The Ecological Society of America, Society for Conservation Biology, The Wildlife Society. *The Wildlife Society Technical Review* 94-2. 112 pp.
- Hoelscher, B. and T.C. Bjornn. 1989. Habitat, density and potential production of trout and char in Pend O'reille Lake tributaries. Project F-71'-R-10, Subproject III, Job No. 8. Idaho Department of Fish and Game, Boise, ID. 72 pp.
- Howell, P.J. and D.V. Buchanan, eds. 1992. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis, OR. 72 pp.
- Howell, P. J., J. B. Dunham, and P. M. Sankovich. 2009. Relationships between water temperatures and upstream migration, cold water refuge use, and spawning of adult bull trout from the Lostine River, Oregon, USA. Published in 2009: *Ecology of Freshwater Fish* 2010:19: 96-106. Malaysia. 11 pp.
- Hudson, J. M., R. Koch, J. Johnson, J. Harris, M. L. Koski, B. Galloway, and J. D. Williamson. 2015. Clackamas River Bull Trout Reintroduction Project, 2014 Annual Report. Oregon Department of Fish and Wildlife and U.S. Fish and Wildlife Service, 33 pp.
- [IDFG] High, B, Meyer, K., Schill, D., and E. Mamer. 2005. Bull trout status review and assessment in the State of Idaho. Grant #F-73-R-27. Idaho Department of Fish and Game. 57pp.
- [IDFG] High, B, Meyer, K., Schill, D., and E. Mamer. 2008. Distribution, abundance, and population trends of bull trout in Idaho. *North American Journal of Fisheries Management* 28:1687-1701.
- IPCC (Intergovernmental Panel on Climate Change). 2007. *Climate change 2007: the physical science basis*. Available: www.ipcc.ch. (February 2007). 1007 pp.

- ISAB (Independent Scientific Advisory Board). 2007. Climate change impacts on Columbia River basin fish and wildlife. ISAB 2007-2. Portland, Oregon. 2007. 146 pp.
- Johnson, L. 1990. State of Nevada, Department of Wildlife, Bull trout management plan. State of Nevada statewide Fisheries Program, project number F-20-26, Job number 2017.4. 17 pp.
- Leary, R.F. and F.W. Allendorf. 1997. Genetic confirmation of sympatric bull trout and Dolly Varden in western Washington. *Transactions of the American Fisheries Society* 126:715-720. 6 pp.
- Leary, R.F., F.W. Allendorf, and S.H. Forbes. 1993. Conservation genetics of bull trout in the Columbia and Klamath River drainages. *Conservation Biology* [CONSERV. BIOL.] 7:856-865.
- Leathe, S.A. and P. Graham. 1982. Flathead Lake Fish Food Habits Study. Environmental Protection Agency, through Steering Committee for the Flathead River Basin Environmental Impact Study. 208 pp.
- Light, J., L. Herger, and M. Robinson. 1996. Upper Klamath basin bull trout conservation strategy, a conceptual framework for recovery. Part one. The Klamath Basin Bull Trout Working Group. 88 pp.
- Magnuson, J.J., Robertson, D.M., Benson, B.J., Wynne, R.H., Livingstone, D.M., Arai, T., Assel, R.A., Barry, R.G., Card, V., Kuusisto, E., Granin, N.G., Prowse, T.D., Stewart, K.M., and Vuglinski, V.S. 2000. Historical trends in lake and river cover in the Northern Hemisphere. *Science* 289:1743-1746. 5 pp.
- Martinez, P. J., P. E. Bigelow, M. A. Deleray, W. A. Fredenberg, B. S. Hansen, N. J. Horner, S. K. Lehr, R. W. Schneidervin, S. A. Tolentino, and A. E. Viola. 2009. Western lake trout woes. *Fisheries* 34:424-442.
- MBTSG (Montana Bull Trout Scientific Group). 1995a. Upper Clark Fork River drainage bull trout status report (including Rock Creek). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 46 pp.
- _____. 1995b. Bitterroot River drainage bull trout status report. Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 34 pp.
- _____. 1995c. Blackfoot River drainage bull trout status report. Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 43 pp.
- _____. 1995d. Flathead River drainage bull trout status report (including Flathead Lake, the North and Middle forks of the Flathead River and the Stillwater and Whitefish River). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 52 pp.
- _____. 1995e. South Fork Flathead River drainage bull trout status report (upstream of Hungry Horse Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 43 pp.

- _____. 1996a. Swan River drainage bull trout status report (including Swan Lake). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 48 pp.
- _____. 1996b. Lower Clark Fork River drainage bull trout status report (Cabinet Gorge Dam to Thompson Falls). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 43 pp.
- _____. 1996c. Middle Clark Fork River drainage bull trout status report (from Thompson Falls to Milltown, including the lower Flathead River to Kerr Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 31 pp.
- _____. 1996d. Lower Kootenai River drainage bull trout status report (below Kootenai Falls). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 39 pp.
- _____. 1996e. Middle Kootenai River drainage bull trout status report (between Kootenai Falls and Libby Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 27 pp.
- _____. 1996f. Upper Kootenai River drainage bull trout status report (including Lake Koocanusa, upstream of Libby Dam). Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 31 pp.
- _____. 1998. The relationship between land management activities and habitat requirements of bull trout. Prepared for Montana Bull Trout Restoration Team. Helena, Montana. 86 pp.
- McIntosh, B.A., J.R. Sedell, J.E. Smith, R.C. Wissmar, S.E. Clarke, G.H. Reeves, and L.A. Brown. 1994. Management history of eastside ecosystems: Changes in fish habitat over 50 years, 1935 to 1992. U.S. Forest Service, Pacific Northwest Research Station, General Technical Report. PNW-GTR 321. 62 pp.
- Meeuwig, M., C. S. Guy, S. T. Kalinowski, and W. Fredenberg. 2010. Landscape influences on genetic differentiation among bull trout populations in a stream-lake network. *Molecular Ecology* 19:3620-3633.
- Minckley, W. L., D. A. Henrickson, and C. E. Bond. 1986. Geography of western North American freshwater fishes: description and relationships to intracontinental tectonism. Pages 519-613 in C. H. Hocutt and E. O. Wiley, editors. *The zoogeography of North American freshwater fishes*. Wiley and Sons, New York.
- McPhail, J.D., and J.S. Baxter. 1996. A Review of Bull Trout (*Salvelinus confluentus*) Life-history and Habitat Use in Relation to Compensation and Improvement Opportunities. University of British Columbia. Fisheries Management Report #104. 37 pp.
- Meehan, W.R. 1991. Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. American Fisheries Society Special Publication 19. 12 pp.

- Meffe, G.K., and C.R. Carroll. 1994. Principles of conservation biology. Sinauer Associates, Inc. Sunderland, Massachusetts. 8 pp.
- Mogen, J. 2013. Bull trout investigations in the Saint Mary River Drainage, Montana – 2010-2012 summary report. U.S. Fish and Wildlife Service Northern Rockies FWCO, Bozeman, Montana.
- Mogen, J. T., and L. R. Kaeding. 2001. Population biology of bull trout (*Salvelinus confluentus*) in the Saint Mary River drainage, progress report 1997-2001. U.S. Fish and Wildlife Service, Bozeman, Montana.
- Mogen, J. T., and L. R. Kaeding. 2005a. Identification and characterization of migratory and nonmigratory bull trout populations in the St. Mary River drainage, Montana. Transactions of the American Fisheries Society 134:841-852.
- Mogen, J. T., and L.R. Kaeding. 2005b. Large-scale, seasonal movements of radiotagged, adult bull trout in the St. Mary River drainage, Montana and Alberta. Northwest Science 79(4):246-253.
- Moore, T. 2006. Distribution and abundance of bull trout and redband trout in Leonard and Deming Creeks, July and August, 2005. Oregon Department of Fish and Wildlife. Corvallis, Oregon.
- Myrick, C.A., F.T. Barrow, J.B. Dunham, B.L. Gamett, G.R. Haas, J.T. Peterson, B. Rieman, L.A. Weber, and A.V. Zale. 2002. Bull trout temperature thresholds:peer review summary. USFWS, Lacey, Washington, September 19, 2002. 14 pp
- NPS (National Park Service). 1992. Value Analysis, Glacier National Park, Divide Creek. West Glacier, Montana.
- Nehlsen, W., J. Williams, and J. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk from California, Oregon, Idaho, and Washington. Fisheries 16(02):4-21. 20 pp.
- Newton, J.A., and S. Pribyl. 1994. Bull trout population summary: Lower Deschutes River subbasin. Oregon Department of Fish and Wildlife, The Dalles, Oregon. Oregon administrative rules, proposed amendments to OAR 340-41-685 and OAR 340-41-026. January 11, 1996. 18 pp.
- ODEQ (Oregon Department of Environmental Quality). 1995. National pollution discharge elimination system permit evaluation report. Facility Bourne Mining Corporation. December 11, 2003. File number 11355. 8pp.
- ODFW (Oregon Department of Fish and Wildlife). 2012. Klamath watershed fish district stock status report, September 2012. ODFW, Klamath Falls, Oregon.
- Porter, M. and M. Nelitz. 2009. A future outlook on the effects of climate change on bull trout (*Salvelinus confluentus*) habitats in the Cariboo-Chilcotin. Prepared by ESSA

- Technologies Ltd. for Fraser Salmon and Watersheds Program, B.C. Ministry of Environment, and Pacific Fisheries Resource Conservation Council. 10 pp.
- Pratt, K.L. 1985. Pend Oreille trout and char life history study. Idaho Department of Fish and Game, Boise, Idaho. 74 pp.
- Pratt, K.L. 1992. A Review of bull trout life history. 00. 5-9. In Proceedings of the Gearhart Mountain Bull Trout Workshop, ed. Howell, P.J. and D.V. Buchanan. Gearhart Mountain, OR. Corvallis, OR: Oregon Chapter of the American Fisheries Society. August 1992. 8 pp.
- Pratt, K.L., and J.E. Huston. 1993. Status of bull trout (*Salvelinus confluentus*) in Lake Pend Oreille and the lower Clark Fork River: (draft report) Prepared for the WWPC, Spokane, WA. 200 pp.
- Quinn, T. P. 2005. The behavior and ecology of pacific salmon and trout. 2005. University of Washington Press. 1st edition. 9 pp.
- Ratliff, D.E., and P.J. Howell. 1992. The status of bull trout populations in Oregon. Pages 10-17 in: P.J. Howell and D.V. Buchanan (eds). Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis. 8 pp.
- Redenbach, Z., and E. B. Taylor. 2002. Evidence for historical introgression along a contact zone between two species of char (Pisces: Salmonidae) in northwestern North America. *Evolution* 56:1021-1035.
- Rich, C.F., Jr. 1996. Influence of abiotic and biotic factors on occurrence of resident bull trout in fragmented habitats, western Montana. MS thesis, Montana State University, Bozeman, MT. 60 pp.
- Rieman, B.E., and J.D. McIntyre. 1993. Demographic and habitat requirements of bull trout *Salvelinus confluentus*. General Technical Report INT-GTR- 302. U.S. Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah. 42 pp.
- Rieman, B.E., and J.D. McIntyre. 1995. Occurrence of bull trout in naturally fragmented habitat patches of varied size. *Transactions of the American Fisheries Society* 124:285-296. 12 pp.
- Rieman, B.E. and J.D. McIntyre. 1996. Spatial and temporal variability in bull trout redd counts. *North American J. of Fisheries Manage.* 16: 132-146. 10pp.
- Rieman, B., and J. Clayton. 1997. Wildfire and native fish: Issues of forest health and conservation of sensitive species. *Fisheries* 22:6-14. 10 pp.
- Rieman, B.E., and J.B. Dunham. 2000. Metapopulations and salmonids: a synthesis of life history patterns and empirical observations. *Ecology of Freshwater Fish* 9:51-64. 14 pp.

- Rieman, B.E., D. Isaak, S. Adams, D. Horan, D. Nagel, C. Luce, D. Myers. 2007. Anticipated Climate Warming Effects on Bull Trout Habitats and Populations Across the Interior Columbia River Basin. *Transactions of the American Fisheries Society*. 136:1552-1565. 16 pp.
- Rode, M. 1990. Bull trout, *Salvelinus confluentus suckley*, in the McCloud River: status and recovery recommendations. Administrative Report Number 90-15. California Department of Fish and Game, Sacramento, California. 44 pp.
- Saunders, D.A., R.J. Hobbs, and C.R. Margules. 1991. Biological consequences of ecosystem fragmentation: A review. *Conservation Biology* 5:18-32. 15 pp.
- Schill, D.J. 1992. River and stream investigations. Job Performance Report, Project F-73-R-13. Idaho Department of Fish and Game, Boise, Idaho. 66 pp.
- Sedell, J.R. and F.H. Everest. 1991. Historic changes in pool habitat for Columbia River Basin salmon under study for TES listing. Draft USDA Report. Pacific Northwest Research Station. Corvallis, OR. 6 pp.
- Sexauer, H.M., and P.W. James. 1997. Microhabitat Use by Juvenile Trout in Four Streams Located in the Eastern Cascades, Washington. Pages 361-370 in W.C. Mackay, M.K. Brown and M. Monita (eds.). *Friends of the Bull Trout Conference Proceedings*. Bull Trout Task Force (Alberta), c/o Trout Unlimited Calgary, Canada. 10 pp.
- Shively, D., C. Allen, T. Alsbury, B. Bergamini, B. Goehring, T. Horning and B. Strobel. 2007. Clackamas River bull trout reintroduction feasibility assessment. Sandy, Oregon, Published by USDA Forest Service, Mt. Hood National Forest for the Clackamas River Bull Trout Working Group.
- Shuter, B.J., and Meisner, J.D. 1992. Tools for assessing the impact of climate change on freshwater fish populations. *GeoJournal* 28(1):7-20. 22 pp.
- Simpson, J.C., and R.L. Wallace. 1982. *Fishes of Idaho*. University Press of Idaho. Moscow, ID. 5 pp.
- Smillie, G. M., and D. Ellerbroek. 1991. Flood hazard evaluation for Divide and Wild creeks, Glacier National Park. Technical Report NPS/NRWRD/NRTR-91/02. Water Resources Division, National Park Service, Fort Collins, Colorado.
- Spruell, P., B.E. Rieman, K.L. Knudsen, F.M. Utter, and F.W. Allendorf. 1999. Genetic population structure within streams: microsatellite analysis of Bull trout populations. *Ecology of Freshwater Fish* 8:114-121. 8 pp.
- Spruell P., A.R. Hemmingsen, P.J. Howell, N. Kandal and F.W. Allendorf. 2003. Conservation genetics of bull trout: Geographic distribution of variation at microsatellite loci. *Conservation Genetics* 4: 17-29. 14 pp.

- Stewart, D.B., N.J. Mochnacz, C.D. Sawatzky, T.J. Carmichael, and J.D. Reist. 2007. Fish life history and habitat use in the Northwest territories: Bull trout (*Salvelinus confluentus*). Canadian Manuscript Report of Fisheries and Aquatic Sciences 2801. Department of Fisheries and Oceans, Winnipeg, MB, Canada, 2007, 54 pp.
- Taylor, B.E., S. Pollard, and D. Louie. 1999. Mitochondrial DNA variation in bull trout (*Salvelinus confluentus*) from northwestern North America: implications for zoogeography and conservation. *Molecular Ecology* 8:1155-1170. 16 pp.
- Thomas, G. 1992. Status of bull trout in Montana. Report prepared for Montana Department of Fish, Wildlife and Parks, Helena, Montana. 108 pp.
- USDA (U.S. Department of Agriculture), and USDI (U.S. Department of the Interior). 1995. Decision Notice/Decision Record Finding of No Significant Impact, Environmental Assessment for the Interim Strategies for Managing Anadromous Fish-producing Watersheds in Eastern Oregon, and Washington, Idaho, and portions of California (PACFISH). 211 pp.
- USFWS (U.S. Fish and Wildlife Service). 1996. Policy Regarding the Recognition of Distinct Vertebrate Population Segments under the Endangered Species Act. *Federal Register* Vol. 61 4722-4725.
- _____. 1998. Determination of threatened status for the Klamath River and Columbia River distinct population segments of bull trout. *Federal Register* Vol. 63 31647-31674. 28 pp.
- _____. 1999. Determination of threatened status for bull trout in the coterminous United States; Final Rule. *Federal Register* Vol. 64 58190-58933. 25 pp.
- _____. 2002a. Bull trout (*Salvelinus confluentus*) draft recovery plan - Chapter 1: Introduction. U.S. Fish and Wildlife Service, Portland, Oregon, October, 2002, 137 pp.
- _____. 2002b. Bull trout (*Salvelinus confluentus*) draft recovery plan - chapter 2 Klamath River. U.S. Fish and Wildlife Service, Portland, Oregon. 93 pp.
- _____. 2004. Draft Recovery Plan for the Coastal-Puget Sound Distinct Population Segment of Bull Trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Portland, Oregon. 297 pp.
- _____. 2008a. Bull trout (*Salvelinus confluentus*) 5-year review: summary and evaluation. Portland, Oregon. 55 pp.
- _____. 2008b. Bull trout draft core area templates - complete core area by core area analysis. W. Fredenberg and J. Chan, editors. U. S. Fish and Wildlife Service. Portland, Oregon. 1,895 pages.
- _____. 2010. Revised designation of critical habitat for bull trout in the coterminous United States. *Federal Register* Vol 75, No. 200. 63898-64070.

- _____. 2015. Recovery plan for the coterminous United States population of bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Portland, Oregon. xii + 179 pp.
- _____. 2015a. Coastal recovery unit implementation plan for bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Lacey, Washington, and Portland, Oregon. 155 pp.
- _____. 2015b. Klamath recovery unit implementation plan for bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Klamath Falls, Oregon. 35 pp.
- _____. 2015c. Mid-Columbia recovery unit implementation plan for bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Portland, Oregon. 345 pp.
- _____. 2015d. Columbia headwaters recovery unit implementation plan for bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Kalispell, Montana, and Spokane, Washington. 179 pp.
- _____. 2015e. Upper Snake recovery unit implementation plan for bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Boise, Idaho. 113 pp.
- _____. 2015f. St. Mary recovery unit implementation plan for bull trout (*Salvelinus confluentus*). U.S. Fish and Wildlife Service, Kalispell, Montana. 30 pp.
- Watson, G., and T.W. Hillman. 1997. Factors affecting the distribution and abundance of bull trout: and investigation at hierarchical scales. *North American Journal of Fisheries Management* 17:237-252. 16 pp.
- WDFW (Washington Department of Fish and Wildlife), FishPro Inc., and Beak Consultants. 1997. Grandy Creek trout hatchery biological assessment. March 1997. Olympia, Washington
- WDFW. 1998. Washington State Salmonid Stock Inventory - Bull Trout/Dolly Vardin. 444 pp.
- WDOE (Washington Department of Ecology). 2002. Evaluating criteria for the protection of freshwater aquatic life in Washington's surface water quality standards - dissolved oxygen: Draft discussion paper and literature summary. Publication Number 00-10-071. Washington Department of Ecology, Olympia, WA, 90 pp.
- Whiteley, A.R., P. Spruell, F.W. Allendorf. 2003. Population Genetics of Boise Basin Bull Trout (*Salvelinus confluentus*). University of Montana, Division of Biological Sciences. Report to the U.S. Forest Service, Rocky Mountain Research Station, Boise, ID. 37 pp.
- Whitesel, T. A., J. Brostrom, T. Cummings, J. Delavergne, W. Fredenberg, H. Schaller, P. Wilson, and G. Zydlewski. 2004. Bull trout recovery planning: a review of the science associated with population structure and size. Science team report #2004-01, U.S. Fish and Wildlife Service, Portland, Oregon. 68 pp.

Wissmar, R., J. Smith, B. McIntosh, H. Li, G. Reeves, and J. Sedell. 1994. A history of resource use and disturbance in riverine basins of eastern Oregon and Washington (early 1800s-1990s). *Northwest Science* 68:1-35. 18 pp.

Ziller, J.S. 1992. Distribution and relative abundance of bull trout in the Sprague River subbasin, Oregon. Pages 18-29 in Howell, P.J. and D.V. Buchanan, editors. *Proceedings of the Gearhart Mountain bull trout workshop*. Oregon Chapter of the American Fisheries Society, Corvallis, OR. 12 pp.

